

DENTAL STATISTICS AND RESEARCH SERIES
Number 40

The National Survey of Adult Oral Health 2004–06

New South Wales

2008

Australian Institute of Health and Welfare
Canberra

Cat. no DEN 176

© Australian Institute of Health and Welfare 2008

This work is copyright. Apart from any use as permitted under the *Copyright Act 1968*, no part may be reproduced without prior written permission from the Australian Institute of Health and Welfare. Requests and enquiries concerning reproduction and rights should be directed to the Head, Media and Communications Unit, Australian Institute of Health and Welfare, GPO Box 570, Canberra ACT 2601.

This publication is part of the Australian Institute of Health and Welfare's Dental statistics and research series. A complete list of the Institute's publications is available from the Institute's website <www.aihw.gov.au>.

ISSN 1321-0254

ISBN 978 1 74024 782 5

Suggested citation

AIHW, Dental Statistics and Research Unit, 2008. The National Survey of Adult Oral Health 2004–06: New South Wales. Cat. no. DEN 176. Dental Statistics and Research Series no. 40. Canberra: Australian Institute of Health and Welfare.

Australian Institute of Health and Welfare

Board Chair

Hon. Peter Collins, AM, QC

Director

Penny Allbon

Published by the Australian Institute of Health and Welfare

Printed by

Contents

Abbreviations.....	v
Symbols.....	vi
Acknowledgments.....	vi
Overview of results.....	1
1 Introduction.....	2
Purpose and organisation of this report	3
Background to the survey	3
Aspects of oral health and dental care relevant to the National Oral Health Plan.....	5
2 Methods	6
Study population and sampling.....	6
Sampled postcodes.....	6
Computer-assisted telephone interview	6
Oral epidemiological examination.....	7
Period of data collection.....	11
Ethical conduct of research	11
Target sample size.....	11
Participation in the survey	12
Data analysis	12
Distribution of sociodemographic and dental access characteristics.....	15
3 Oral health status	16
Complete tooth loss	16
Inadequate natural dentition among dentate people.....	18
Denture wearing by dentate people	20
Average number of teeth per person missing due to pathology.....	22
Prevalence of untreated coronal decay	24
Percentage of people with untreated root decay	26
Percentage of people with one or more filled teeth.....	28
Average number of decayed, missing and filled teeth per person	30
Prevalence of moderate or severe periodontitis	32
Prevalence of deep pocket depth	34
Prevalence of 4+ mm clinical attachment loss.....	36
Prevalence of gingival inflammation.....	38

4	Oral health care	40
	Dental attendance within the preceding 12 months.....	40
	Attendance at private dental practice	42
	Payments by patients for dental care	44
	Government-subsidised dental care in private sector	46
	People’s usual pattern of dental visits.....	48
	Usual attendance at the same dentist.....	50
	Usual dental attendance for a check-up.....	52
	Dental care avoided or delayed due to cost	54
	Recommended dental treatment foregone due to cost	56
	Difficulty paying a \$100 dental bill.....	58
	Percentage of people avoiding foods due to dental problems	60
5	Oral health perceptions	62
	Percentage of people rating their oral health as fair or poor	62
	Percentage of people experiencing toothache.....	64
	Percentage of people experiencing orofacial pain.....	66
	Perceived need for dentures	68
	Perceived need for dental extraction or filling.....	70
	Perceived need for a dental check-up	72
	Perceived urgency of dental treatment needs.....	74
	Age-standardised comparison between government health cardholders and non-health cardholders	76
	Age-standardised comparison between the dentally insured and the uninsured.....	78
	Appendix	80
	Sample counts	80
	Glossary	82
	References	85
	List of tables	88

Abbreviations

AAP	American Academy of Periodontology
AHMAC	Australian Health Ministers' Advisory Council
AIHW	Australian Institute of Health and Welfare
ARCPOH	Australian Research Centre for Population Oral Health
CAL	clinical attachment loss
CATI	computer-assisted telephone interview
CDC	US Centers for Disease Control and Prevention
CEJ	cemento-enamel junction
DMFT	number of decayed, missing and filled permanent teeth
DSRU	Dental Statistics and Research Unit
IRSAD	Index of Relative Socioeconomic Advantage/Disadvantage
NHANES	US National Health and Nutrition Examination Survey
NHMRC	National Health and Medical Research Council
NOHSA	National Oral Health Survey of Australia
NSAOH	National Survey of Adult Oral Health
SEIFA	Socioeconomic Indices for Areas

Place abbreviations

ACT	Australian Capital Territory
NSW	New South Wales
NT	Northern Territory
Qld	Queensland
SA	South Australia
Tas	Tasmania
UK	United Kingdom
US	United States
Vic	Victoria
WA	Western Australia

Symbols

\$	Australian dollars
%	per cent
..	not applicable
–	nil
>	greater than
<	less than
≥	greater than or equal to
≤	less than or equal to
<0	estimate is less than zero

Acknowledgments

This publication was prepared by the following authors at the Australian Research Centre for Population Oral Health (ARCPOH):

Authors

Loc Do

Anne Ellershaw

Jane Harford

Liana Luzzi

Kaye Roberts-Thomson

Gary Slade

John Spencer

Editorial team

Alison McLean and Lorna Lucas (ARCPOH) contributed to the editing of this publication and proofreading was by Jo Mason. All three worked to improve the consistency, layout and readability of the text.

NSW Survey personnel

Survey managers	Shanti Sivaneswaren
Appointment coordinators	Mr Mitchell Flynn, Ms Kawkab (Kazzie) El-Souss, Mr Gary Law
Dentist examiners	Dr Sue Buchanan, Dr Maria Chmielowie, Dr Stephen Chui, Dr Ilana Fisher, Dr Christopher Harle, Dr Thomas Hasson, Dr Peter Hill, Dr Kartusha Hull, Dr Paul Kotala, Dr Emily Kyaw, Dr Zhen Mashen
Dental recorders	Ms Muna Aldalo, Ms Veronica Erdes, Ms Lea Hope, Ms Lexie McLaren, Ms Judy Miekle, Ms Rachael Moreau, Ms Sally Newall, Ms Rowena Palogan, Ms Julie Plimmer, Ms Sudha Raju, Ms Michelle Woolfe

Scientific Advisory Committee members

Dr Peter Barnard, Australian Dental Association
Professor Mike Morgan, The University of Melbourne
Mr Gary Niedorfer, Australian Bureau of Statistics
Professor Brian Oldenburg, Monash University
Professor Eric Reynolds, The University of Melbourne
Dr Ken Tallis, Australian Institute of Health and Welfare

Community and Professional Advisory Committee members

Dr Louise Brown, Australian and New Zealand Academy of Periodontists
Ms Samantha Edmonds, Council of Social Service of NSW (NCOSS)
Mr Errol Evans, Oral Health Unit, Queensland Health (alternate Dr Paul Wood)
Ms Chris Morris, South Australian Dental Service
Dr David Neesham, Dental Health Services, Western Australia (alternate Dr Martin Glick)
Dr Bill O'Reilly, Australian Dental Association Federal Executive
Ms Lindsay Simmons, Council of the Ageing
Ms Tracey Slater, Department of Human Services, Victoria

Funding sources for the 2004–06 National Survey of Adult Oral Health

National Health and Medical Research Council, Project Grant #299060
National Health and Medical Research Council, Project Grant #349514
National Health and Medical Research Council, Capacity Building Grant #349537
Australian Government Department of Health and Ageing, Population Health Division
Australian Institute of Health and Welfare
Colgate Oral Care
Australian Dental Association
US Centers for Disease Control and Prevention, Research Participation Program

ARCPOH interviewers

CATI supervisors

Scott Crowley, Meredith Morgan

CATI interviewers

Jan Anderson, Olivia Baker, Jenny Cates, Debra Flack,
Susie Garner, Alex Holland, Jessica Holder, Dora Jankumas,
Claire Leske, Peter Monaghan, Patricia Moyle, Grant Noble,
Alison Packer, Mary Porter, Jenny Proeve, Verity Proeve,
Edmund Redgrove, Joanne Rogers, Joy Simmons,
Kaye Sparrow, Lachlan Spencer, Michelle Storan, Sue Tanner,
Zoe Taylor, Theodora Trovas, Jenny Venn, Tammy White

Overview of results

This report describes levels of oral health in the adult population of New South Wales (NSW) at the beginning of the twenty-first century. The findings are from the 2004–06 National Survey of Adult Oral Health (NSAOH). In NSW, 3,630 people were interviewed and 1,099 people were dentally examined for the survey. This report presents percentages and means for 30 oral health indicators in tables that compare three age groups and classify people according to five sociodemographic characteristics: sex, residential locality, socioeconomic status of residential postcode, government health card status and dental insurance status.

Oral health status

- 5.5% of people had no natural teeth and among dentate people, an average of 4.9 teeth per person were missing. These and two other indicators of tooth loss were associated with residential location, socioeconomic status, government health card status and dental insurance status.
- 27.1% of people had untreated dental decay and an average of 12.8 teeth per person were decayed, missing or filled. Prevalence of those conditions was associated with the same four sociodemographic characteristics.
- 17.2% of people had inflamed gums and 23.0% had moderate or severe gum disease. Two indicators of gum disease occurred more frequently among government health cardholders than non-cardholders.

Oral health care

- 60.3% of people had visited a dentist within the preceding 12 months and 54.9% said they usually did so. These and two other measures of dental attendance varied according to residential location, socioeconomic status, government health card status and dental insurance status.
- 81.4% of people had a dentist that they usually attended, although 29.8% said that they avoided or delayed dental care due to its cost. Barriers to dental care were associated with all five sociodemographic characteristics.

Oral health perceptions

- 19.7% of people said they had avoided some foods due to dental problems, and 16.2% had experienced toothache, in the preceding 12 months. Perceptions of poor oral health were associated with all five sociodemographic characteristics.
- 31.1% of people felt they needed an extraction or filling, although only 7.1% said they needed dentures. Perceived dental treatment needs were associated with socioeconomic status, government health card status and dental insurance status.

Age-standardised analysis revealed that government health cardholders had poorer outcomes for 23 of the 29 indicators, while the uninsured had poorer outcomes for 23 of the 30 indicators reported.

1 Introduction

This report presents findings from the NSW component of the 2004–06 National Survey of Adult Oral Health (NSAOH). Information was collected using interviews and standardised dental examinations that were conducted among a random sample of NSW residents aged 15 years or more. Three major themes are reported in chapters describing oral health status, oral health care and perceptions of oral health. Statistics summarising those themes are tabulated for the NSW adult population and for three age groups that are further classified according to: sex, residential locality, socioeconomic status of the area in which they live, government health cardholder status and dental insurance.

The 2004–06 NSAOH took place 17 years after the first oral examination survey of Australians conducted in the six states and the Australian Capital Territory (Barnard 1993). State/territory reports from that 1987–88 National Oral Health Survey of Australia (NOHSA) highlighted variations among age groups, between the sexes and between people living in or outside capital cities. The major findings reported from the survey were:

- children’s dental decay rates were low by historical standards and when compared internationally
- nearly one-half (48%) of adults had made a dental visit within the preceding year, the majority of them to a private dental practice (88%)
- however, 44% of adults were found to need one or more dental fillings
- the percentage of Australians with complete tooth loss had reduced compared with earlier interview surveys, although 50% of people aged 65 years or more had no natural teeth and
- one of the four national oral health targets had been achieved, and it was expected that the remaining three targets would be achieved by 2000.

However, the first survey did not collect information about government health cardholder status or socioeconomic status, and results were not contrasted between insured and uninsured.

In the 17-year period since the NOHSA, there has been substantial growth in public sector dental care and dental insurance. Increasingly, national and state/territory health goals call for reductions in socioeconomic inequalities in health, including oral health. For those reasons, this report includes a focus on the relationship between oral health and indicators of socioeconomic status and access to dental care, as well as the traditional demographic markers of age, sex and residential location.

Purpose and organisation of this report

The purpose of this report is to provide a descriptive ‘snapshot’ of oral health in the adult population of NSW. The findings are intended to provide up-to-date evidence that can contribute to the development of oral health policies and programs in NSW.

This introductory chapter outlines the motives for undertaking the survey. Chapter 2 reviews the survey’s methods and describes the population distribution of sociodemographic and dental access characteristics presented in later tables. Statistical findings regarding oral health status are tabulated and described in Chapter 3, followed by statistical findings regarding oral health care (Chapter 4) and perceptions of oral health (Chapter 5). The Appendix contains additional tables of oral health statistics for conventional 10-year age groups. These are narrower than the age ranges reported in the chapters, and are presented to permit comparisons with surveys conducted at other places and other times.

The national report of the survey’s findings (Slade et al. 2007) provides additional details about the survey, including participation rates and analysis of potential biases due to non-participation. The national report also presents qualitative findings from ‘oral histories’ conducted with a small number of survey participants to document historical influences on the nation’s oral health. Further appendix material is available at:

<<http://www.arcpho.adelaide.edu.au/project/distribution/NSAOH.html>>.

Background to the survey

Up-to-date information about population oral health is important because oral diseases have broad implications for the health of the public. Dental problems are ranked among the most frequently reported illness episodes by Australians (AIHW 2000) and provision of dental care accounts for 6.6% of recurrent health expenditure in 2005–06 (AIHW 2007). In the United States the Surgeon General characterised oral disease as a ‘silent epidemic’ (Surgeon General 2000).

In the 17 years following the 1987–88 NOHSA, no state-wide oral examination surveys of adults have been conducted. Instead, published oral examination surveys were restricted to special groups of the adult population and often they were conducted within selected locations in states. They included studies of oral health in:

- military recruits (Dawson & Smales 1994; Hopcraft & Morgan 2003a,b, 2005, 2006; Morgan et al. 1992)
- adults in Melbourne (Wright et al. 1994)
- community dwelling elderly people (Bergman et al. 1991; Chalmers, Carter & Spencer 2002; Slade et al. 1993; Slade & Spencer 1995,1997; Thomson et al. 1995)
- elderly people living in nursing homes or hostels (Chalmers, Carter, Fuss et al. 2002; Chalmers, Hodge et al. 2002; Chalmers et al. 2005; Saub & Evans 2001)
- Aboriginals and Torres Strait Islanders (Endean et al. 2004; Smith et al. 2007)
- immigrants (Marino et al. 2001, 2007) or refugees (Kingsford Smith & Szuster 2000)
- prisoners (Osborn et al. 2003)
- patients receiving dental care in public dental services (Brennan et al. 2000, 2001, 2007; Brennan & Spencer 2004) and
- patients with selected medical conditions (Coates et al. 1996, 2000).

By the late 1990s, several collaborative efforts among federal and state/territory stakeholders attempted to secure support for a second national oral health survey, although none were funded. Renewed impetus for a national survey began with the work of the National Advisory Committee on Oral Health (AHMAC 2001). The committee formulated a National Oral Health Plan for the period 2004–13 comprising seven action areas:

- promotion of oral health across the population
- children and adolescents
- older people
- people with low income and social disadvantage
- people with special needs
- Aboriginal and Torres Strait Islander people and
- workforce development.

One of four-short term goals listed for the plan's first action area was the conduct of a national survey of adult oral health. Fulfilment of that goal became possible in 2003 when researchers at the Australian Research Centre for Population Oral Health (ARCPOH) in The University of Adelaide sought project grant funding from the National Health and Medical Research Council (NHMRC). The proposal was for funding to support a collaborative project that pooled resources already committed or promised from the following sources: funding from the Australian Government Department of Health and Ageing to the Dental Statistics and Research Unit (DSRU) within ARCPOH to undertake a telephone interview survey; commitment of staff from oral health sections within state and territory health departments to conduct oral epidemiological examinations; and core funding from the Australian Institute of Health and Welfare (AIHW) to DSRU. Following peer review, the NHMRC awarded a project grant to ARCPOH in November 2003.

Aspects of oral health and dental care relevant to the National Oral Health Plan

The National Oral Health Plan outlined nine population indicators that were informative in developing the plan and that are cited as key performance indicators to evaluate the outcomes of the plan. This survey reports findings that relate to six of those key performance indicators:

- The percentage of the dentate population reporting a social impact (for example toothache, difficulty chewing, concerned about appearance) because of problems with teeth, mouth or gums in the last 12 months, by age group, living circumstance, government health cardholder status, Indigenous identity and special needs.
- The percentage of the population with untreated decay, by age group, living circumstance, government health cardholder status and Indigenous identity.
- The proportion of the dentate population with a maximum periodontal pocketing of 3.5 mm and 5.5 mm, by age group.
- The mean number of missing teeth and proportion of existing teeth with untreated decay, by age group, living circumstance, government health cardholder status and Indigenous identity.
- The percentage of the dentate population who visited a dental practitioner in the last 2 years, by age group, living circumstance, government health cardholder status and Indigenous identity.
- The percentage of the dentate population whose reason for visiting a dental practitioner in the last 12 months was for a check-up, by age group, living circumstance, government health cardholder status and Indigenous identity.

2 Methods

Full details of the survey's methods have been described in Chapter 2 of the national report (Slade et al. 2007). The following summary highlights the main methodological features of the survey.

Study population and sampling

A three-stage, stratified clustered sampling design was used to select people from the target population of Australian residents aged 15 years or more:

- Postcodes were sampled at random from capital city and non-capital city strata in six states and the Northern Territory, and from a single stratum in the Australian Capital Territory. Postcodes represented the geographic clustering in the design and were selected with probability proportional to size, where size was defined as the number of households listed in the 'electronic white pages' in each postcode.
- A systematic sample of households listed in the 'electronic white pages' was selected for each sampled postcode. Thirty households per metropolitan stratum and 40 households per ex-metropolitan stratum were selected.
- One person aged 15 years or more was randomly selected per household. In households with only one person aged 15 years or more, that person was selected. In other households telephone interviewers asked for the name of the person aged ≥ 15 years who most recently had had a birthday and the name of the person aged ≥ 15 years who would next have a birthday. A computer algorithm then selected one of those two people at random.

Sampled postcodes

In NSW the following postcodes were sampled: 2008, 2011, 2018, 2022, 2026, 2030, 2032, 2035, 2037, 2040, 2044, 2048, 2062, 2066, 2068, 2072, 2075, 2077, 2083, 2087, 2089, 2094, 2097, 2100, 2106, 2112, 2114, 2118, 2121, 2126, 2132, 2137, 2142, 2145, 2146, 2148, 2151, 2153, 2155, 2161, 2164, 2166, 2170, 2176, 2192, 2195, 2200, 2204, 2207, 2210, 2213, 2218, 2221, 2225, 2229, 2230, 2233, 2250, 2251, 2257, 2259, 2261, 2262, 2280, 2284, 2289, 2299, 2318, 2320, 2326, 2333, 2350, 2360, 2402, 2429, 2444, 2450, 2460, 2478, 2482, 2500, 2517, 2527, 2535, 2540, 2546, 2560, 2564, 2567, 2576, 2587, 2620, 2641, 2650, 2666, 2738, 2747, 2750, 2759, 2761, 2767, 2770, 2780, 2790, 2800, 2850, 2880.

Computer-assisted telephone interview

Self-reported information about oral health and characteristics associated with it was obtained through telephone interviews. Interviewers read questions from a computer screen and recorded answers directly onto the computer. They were conducted from a dedicated computer-assisted telephone interview (CATI) suite at University of Adelaide research offices. The methods were based on those advocated by Dillman (2000), including the mailing of a letter to households prior to telephoning, a protocol for contacting each household, and standardised procedures for asking questions and recording answers.

Interviews were conducted by 29 interviewers, each of whom was trained in the Survey methods. Every effort was made to interview the target person although, in certain circumstances, the questions were answered by another adult in the form of a proxy interview.

The interview consisted of 79 questions, several with multiple response categories. A copy of the questions used is included in an Appendix available online:

<<http://www.arcpoh.adelaide.edu.au/project/distribution/NSAOH.html>>.

Oral epidemiological examination

Information about clinical oral status was collected during standardised dental examinations conducted by dentists who undertook training in the survey procedures. Examinations were limited to people who reported having some or all of their own natural teeth at the time of the interview. Examining dentists followed a standardised protocol to record levels of tooth loss, dental decay experience, tooth wear and – for subjects with no medical contraindications to periodontal probing – signs of gum disease. During data collection, replicate examinations were conducted for approximately five study participants per examiner to evaluate the consistency of their findings when judged against the principal survey examiner.

There were 30 examiners nationwide (Table 1). Prior to their work on the survey, they undertook a 2-day training and calibration session at The University of Adelaide. Separate training sessions were held for the examination teams from each state and territory. Prior to the scheduled training session, each examiner was sent a 50-page manual and a DVD detailing the survey protocol, including the criteria and coding for the examination.

Table 1: Distribution of examiners and examinations among states and territories

State	No. of examiners	No. of people examined	No. of examinations per examiner		
			Minimum	Maximum	Mean
NSW	11	1,113	32	164	101
Vic	3	1,181	267	585	394
Qld	3	824	217	305	275
SA	2	629	241	388	315
WA	3	470	134	196	157
Tas	3	385	49	186	128
ACT	2	386	125	261	193
NT	3	517	154	203	172
All states	30	5,505	32	585	184

Scope of examination

Survey participants were examined in a supine position in standard dental chairs with illumination provided by the chair's overhead dental light. Examiners used an intra-oral mirror that additionally had its own battery-powered light source. A periodontal probe with 2-mm markings was used to record distances, for example when assessing periodontal destruction (described further below); however, sharp explorers were not used and no radiographs were taken. Full details of the examination protocol are provided online:

<<http://www.arcpoh.adelaide.edu.au/project/distribution/NSAOH.html>>.

The following overview summarises criteria used to assess the main oral health variables reported in this volume.

Tooth loss

For people aged less than 45 years, examiners distinguished between missing teeth that had been extracted due to decay or periodontal disease and teeth that were absent for any other reason (that is, congenitally missing; unerupted; or extracted for orthodontics, trauma or impaction). For people aged 45 years or more, no such distinction was made, so that an extracted or otherwise absent tooth was recorded as missing. Dental implants, root fragments and deciduous teeth were coded separately and not counted as missing or absent teeth.

Replacement teeth

All lost teeth were further classified as replaced or not replaced by a fixed bridge or a removable denture that was worn to the examination.

Decay experience of coronal tooth surfaces

All teeth present were subdivided into five tooth surfaces: mesial, buccal, distal, lingual, and either occlusal (for premolars or molars) or incisal (for incisors and canines). Each coronal surface was assessed and categorised using visual criteria (no explorer was used) and one of the following codes was assigned:

- decay: cavitation of enamel or dentinal involvement or both are present
- recurrent caries: visible caries that is contiguous with a restoration
- filled unsatisfactorily: a filling placed for any reason in a surface that requires replacement but that has none of the above conditions
- filling to treat decay: a filling placed to treat decay in a surface that had none of the above conditions
- filling placed for reasons other than decay: in a surface that has none of the above conditions (incisors and canines only)
- fissure sealant: where none of the above conditions were found
- sound: when none of the above conditions was found.

Decay experience of tooth root surfaces

All teeth present were subdivided into four root surfaces: medial, buccal, distal and lingual. Each root surface was assessed visually and, if necessary, using a ball-ended periodontal probe. One of the following codes was assigned:

- decay: a discrete, well-defined or discoloured lesion on the root surface that is soft to exploration using the periodontal probe
- recurrent caries: detectable caries that is contiguous with a restoration
- filled unsatisfactorily: a filling placed for any reason in a surface that has unacceptable defects but meeting none of the above conditions
- filled root surface: one or more permanent restorations placed for any reason but meeting none of the above conditions
- wear of 2 mm or more: recorded only on buccal surfaces with none of the above conditions
- sound root surface: when none of the above conditions was found
- no visible root surface.

Periodontal tissue destruction

The assessment of periodontal tissue destruction was based on methods used in the US National Health and Nutrition Examination Survey (NHANES 2005). Assessments were made of probing pocket depth and gingival recession, both recorded in millimetres using a periodontal probe that had 2-mm markings. Measurements were made at the mesio-buccal, mid-buccal and disto-buccal aspects of all teeth present, except for third molars. All fractional millimetre measurements were rounded down to the lowest whole millimetre before calling the number. For recession, the cemento-enamel junction (CEJ) was identified or its position was estimated (for example, if a filling obscured its position), and the distance from the CEJ to the free gingival margin was recorded in millimetres. When the CEJ was subgingival, the number called was negative; otherwise it was positive. For probing pocket depth, the distance from the free gingival margin to the bottom of the periodontal crevice/pocket was called.

Examiners did not make a direct measurement of clinical attachment loss; instead, it was computed during data analysis.

Gingival inflammation around six index teeth

The Loe and Silness (1963) gingival index was used to assess inflammation of the marginal gingival tissues around six index teeth (if present) – the most anterior molar in each dental quadrant (up to four teeth), the right maxillary central incisor and the left mandibular central incisor. Pressure was applied to the free gingival margin on the buccal aspect of the tooth by swiping with the side of a periodontal probe that was held at approximately 90 degrees to the long axis of the tooth. One of the following codes was assigned:

- severe inflammation: marked redness and oedema, ulceration or tendency to spontaneous bleeding
- moderate inflammation: redness, oedema, glazing or bleeding after applying pressure with the probe
- mild inflammation: slight change in colour or slight oedema but no bleeding after applying pressure with the probe
- none of the above.

Data recording for examinations

Each code called by an examiner was recorded directly onto a laptop computer by state/territory staff who had experience in clinical dental procedures. They were trained in use of the software during the 2-day training session for examination teams held at The University of Adelaide.

Assessment of inter-examiner reliability

In order to measure inter-examiner reliability, the principal survey examiner attended examination sessions for all but one examiner to conduct masked replicate examinations of survey participants. The remaining examiner withdrew from the survey after completing 32 examinations. Replicate examination entailed assessments of tooth presence, periodontal assessment of teeth in one jaw, and assessment of caries experience in both crowns and roots of teeth. The observed levels of agreement for most oral health indicators were equivalent to benchmarks reported for national oral health surveys conducted in the United Kingdom and the United States.

Period of data collection

Data collection began in July 2004 and was completed in September 2006 (Table 2). Interviews were timed to begin approximately 1 month prior to the planned start of examinations in each jurisdiction.

Table 2: Periods of data collection in states and territories

State/territory	Dates of interviews		Dates of examinations	
	Beginning	End	Beginning	End
ACT	July 2004	October 2004	July 2004	October 2004
SA	September 2004	December 2004	September 2004	May 2005
WA	October 2004	March 2005	November 2004	May 2005
Vic	January 2005	September 2005	February 2005	September 2005
NSW	May 2005	November 2005	June 2005	July 2006
NT	August 2005	October 2005	September 2005	March 2006
Tas	January 2006	May 2006	March 2006	September 2006
Qld	March 2006	September 2006	June 2006	September 2006
Australia	July 2004	September 2006	July 2004	September 2006

Ethical conduct of research

This project was reviewed and approved by The University of Adelaide's Human Research Ethics Committee. Interviewed subjects provided verbal consent prior to answering questions. All examined subjects provided signed, informed consent prior to the examination.

Target sample size

Sample size requirements were calculated for a range of key outcome variables to be reported nationally. One outcome, the capacity to detect a 25% or greater reduction in national age-specific estimates of mean number of decayed teeth since 1987–88, was nominated as the critical threshold that should be detectable with standard statistical power of 80%. Another outcome was a capacity to detect a 10% or greater reduction in national age-specific mean DMFT. This identified a need for 7,500 examinations and 13,560 interviews, assuming a 65% participation rate in the examination. The sample size within each state and territory was planned to be approximately proportional to the population of the jurisdiction.

Participation in the survey

National participation rates were lower than intended, both in the interview, where 49.0% of sampled people participated, and the examination, where 43.7% of those eligible took part. Interview participation rates varied from 43.9% in NSW to 61.8% in SA. Examination rates varied from 33.2% in NSW to 57.5% in SA (Table 3).

Table 3: Number and percentage of people sampled, interviewed and examined^(a)

	No. of people sampled	No. of people interviewed	Per cent of sampled people interviewed	No. of people eligible for exam	No. of people examined	Per cent of eligible people examined
Australia	28,812	14,123	49.0	12,606	5,505	43.7
State/territory						
NSW	8,270	3,630	43.9	3,310	1,099	33.2
Vic	6,013	2,667	44.4	2,360	1,181	50.0
Qld	4,219	2,052	48.6	1,841	824	44.8
SA	2,159	1,335	61.8	1,093	629	57.5
WA	2,365	1,290	54.5	1,109	470	42.4
Tas	1,745	1,042	59.7	873	385	44.1
ACT	1,892	1,025	54.2	981	400	40.8
NT	2,149	1,082	50.3	1,039	517	49.8

(a) Unweighted data.

Data analysis

The aim of the data analysis was to generate summary statistics describing oral health for the NSW population. With the exception of data regarding participation rates, results in this report have been weighted to compensate for individuals' different probabilities of selection and survey participation rates. For the telephone interview survey, weights were adjusted to ensure survey estimates were consistent with the 2005 Australian Bureau of Statistics Estimated Residential Population data. For the oral examination survey, which was restricted to dentate people aged 15 years or more, estimates of the dentate population were derived from the telephone interview survey and used to derive examination weights. This means that results can be generalised to the NSW population.

Tables 35 and 36 contain age-standardised estimates for each indicator presented in preceding tables. Age standardisation is a statistical procedure that aims to remove any effects of age that might account for differences in each oral health indicator between the two comparison groups: health cardholders versus non-health cardholders (Table 35) and insured versus non-insured people (Table 36). For these tables, percentages and means were standardised using the direct method. The reference population was the 2005 Australian Estimated Residential Population classified into 14 five-year age categories within the range 15–84 years and a fifteenth category aged 85 years of more.

Presentation of results

Oral health measures are tabulated for each of three age groups representing the survey participant's age reported in the telephone interview, plus an 'all ages' summary. The three age groups are 15–34 years, 35–54 years and ≥55 years. The tables report estimates for mutually exclusive subgroups of people created for each of six characteristics based on responses to the telephone interview questions. The subgroups and unweighted number of respondents are listed in the Appendix to this volume and the six characteristics are described below:

Sex was classified as 'Male' or 'Female' recorded during the interview.

Residential location was classified as 'Capital city' or 'Other places' based on the sampling postcode used in selection of households.

Postcode socioeconomic status was used to classify individuals according to the Index of Relative Socioeconomic Advantage/Disadvantage (IRSAD) of the postcode in which they lived. The IRSAD is an aggregate measure of a postcode's socioeconomic status based on characteristics of its residents recorded in the 2001 Population Census. A postcode that has a relatively high proportion of people with high incomes or a skilled labour force is assigned a relatively higher value on this index. Conversely, a low score on the index indicates that an area has a higher proportion of individuals with low incomes and more people who work in unskilled occupations. Postcodes were classified into three groups of ascending socioeconomic status, each group comprising approximately one-third of the NSW population. This type of analysis is said to be 'ecological' because it is not based on individuals own socioeconomic status, but on the socioeconomic status of the area in which they live. Hence, care should be taken in the interpretation of results – because Socioeconomic Indices for Areas (SEIFA) scores refer to areas, not individuals, results are not interpretable at the level of the individual.

Government health card status identified whether or not people were covered either by a pensioner concession card or health care card. Both cards are issued according to a means test administered by Centrelink, an agency of the Australian Government's Family Assistance Office. People with either card and their dependents are eligible for public-sector dental care in most states and territories.

Place of last dental visit further disaggregated health cardholders according to the location of their last dental visit. The latter was established during the interview by asking people 'Where did you make your last dental visit?'. Health cardholders who responded 'Government dental clinic' or 'School dental service' were classified as 'Cardholder/Public'. Otherwise, eligible people were classified as 'Cardholder/Non-public' if they reported any of the other locations: Private dental practice (including specialist); Dental technician; Clinic operated by health insurance fund; Armed Services/Defence Force clinic; Other site. People who were not health cardholders were classified as 'Non-cardholder/Non-public' regardless of their reported visit location.

Dental insurance coverage was based on responses to the question 'Do you have private insurance cover for dental expenses?'. People were classified as insured if they responded 'yes' and uninsured if they responded 'no'.

Criteria for determining statistical significance

As with any survey where data are collected from only some of the people in the population, proportions and means in this report are estimates of the true population values. The estimates have some degree of uncertainty, which is expressed in this report using 95% confidence intervals (95% CIs). The 95% CI signifies the likely lower and upper limits of the range of values within which the true population percentage would fall. In this context 'likely' means that there is a 95% probability that the true population value lies between those two values.

In this report 95% CIs are used additionally as a guideline to identify differences between population subgroups that are statistically significant. Specifically, when there is no overlap between 95% CIs for two groups, the difference between the groups is deemed to be statistically significant. This criterion for judging statistical significance is more conservative than the alternative method of calculating P-values. In fact, when 95% CIs do not overlap, it means that a test of statistical significance for the difference between the groups would have a P-value of less than 0.05 (the conventional threshold used in many reports), and it could be as small as less than 0.005. The 'conservative' nature of the criterion used in this report comes about because 95% CIs that overlap to a small degree could, nevertheless, be found to differ to a statistically significant degree (at $P < 0.05$) using a hypothesis test.

Data files were managed and summary variables computed using SAS software version 9.1.¹ Means and their associated 95% CIs were generated using SUDAAN software release 9.0.0.² The SUDAAN procedures used sampling weights to generate population estimates and calculated 95% CIs that allowed for the complex sampling design used in this survey. To do so, 'with replacement' sampling was specified with two levels of stratification (state and section of state). The subject's sampling postcode was specified as the primary sampling unit, which was used by SUDAAN as the clustering variable.

¹ SAS Institute Inc. 100 SAS Campus Drive, Cary, NC 27513-2414, USA.

² Research Triangle Institute. PO Box 12194, Research Triangle Park, NC 27709-2194, USA.

Distribution of sociodemographic and dental access characteristics

Approximately one-half of the NSW population was female, with little variation in the proportion among age groups (Table 4). Two-thirds lived in the capital city, a proportion that was lower in progressively older age groups. By design, people of all ages were approximately evenly distributed among tertiles of postcode, socioeconomic status. However, older people were more likely than younger people to live in areas with lower socioeconomic status. Approximately one-quarter of the population were government health cardholders, although the proportion was noticeably greater for people aged 55 years or more. However, government health cardholders were less likely to have last attended a public dental clinic than other dental care providers, a pattern that was consistent in each age group. Nearly one-half of the NSW population had dental insurance, a figure that did not vary substantially among age groups.

Table 4: Percentage of people with selected sociodemographic and dental access characteristics in the NSW population and three age groups

	All ages	Age group (years)		
		15–34	35–54	>=55
Sex				
Males	50.2	50.6	50.1	49.5
Females	49.8	49.4	49.9	50.5
Residential location				
Capital city	64.1	67.6	63.9	59.6
Other places	35.9	32.4	36.1	40.4
Postcode socioeconomic status				
Lowest	30.3	27.2	32.6	31.5
Middle	32.3	32.0	32.8	32.0
Highest	37.4	40.8	34.6	36.5
Government health card				
Health care card or pensioner concession card	26.4	20.0	17.4	47.2
Neither card	73.6	80.0	82.6	52.8
Place of last dental visit				
Cardholder/Public	8.0	7.0	7.3	10.4
Cardholder/Non-public	18.4	13.0	10.1	36.8
Non-cardholder/Non-public	73.6	80.0	82.6	52.8
Dental insurance				
Insured	46.6	47.8	46.6	45.0
Uninsured	53.4	52.2	53.4	55.0

3 Oral health status

Complete tooth loss

In NSAOH, complete tooth loss was assessed in the interview by asking people 'Do you have any of your own natural teeth?'. People who answered 'no' were classified as edentulous. In NSW, edentulous people represented 5.5% of the population aged 15 years or more (Table 5), which was slightly lower than the national estimate of 6.4% (Slade et al. 2007).

Key findings

- The prevalence of edentulism was strongly associated with age, being negligible among 15–34-year-olds but affecting 16.0% of NSW adults aged 55 years or more.
- Females were almost twice as likely as males to be edentulous, both for all ages combined and for people aged 55 years or more.
- People living outside the capital city were almost twice as likely as those living in Sydney to be edentulous, both for all ages combined and for people aged 55 years or more.
- Prevalence was more than twice as high among people living in postcodes with low socioeconomic status than in postcodes with high socioeconomic status. However, there was no statistically significant difference in prevalence between middle and low socioeconomic postcodes.
- Government health cardholder status was associated with some of the largest differences observed between population groups in prevalence of edentulism. For example, among people aged 55 years or more, prevalence was three-fold greater among government health cardholders compared with non-government health cardholders.
- Among the population of government health cardholders, there was a tendency for greater prevalence in people whose last dental visit was to the public sector than those who attended elsewhere. However, 95% CIs were large in these groups, with the consequence that differences were not statistically significant.
- People with no dental insurance were also much more likely than the insured to be edentulous.

Discussion

As emphasised in the national report, variation among age groups in prevalence of edentulism can be attributed primarily to the differing historical experiences of generations born in different time periods during the 20th century, rather than to the effects of ageing. Because edentulism prevalence was so strongly dependent upon age group, comparisons between population groups were observed most clearly for the oldest age group. In contrast, among 35–54-year-olds, 95% CIs for estimated prevalence frequently overlapped between population groups (for example, when comparing males and females). Among 15–34-year-olds, prevalence estimates were all zero and therefore did not reveal any differences between population groups.

In summary, complete tooth loss in NSW was a condition observed infrequently below the age of 55 years, while among people aged 55 years or more, it was most likely to occur in socioeconomically disadvantaged groups.

Table 5: Percentage of adults with complete tooth loss

		Population: all people Age (years)			
		All ages	15–34	35–54	≥55
All people	Per cent of people	5.5	0.0	1.7	16.0
	<i>95% CI</i>	4.8–6.3	—	1.1–2.9	13.9–18.4
Sex					
Males	% of people	3.9	0.0	1.3	11.9
	<i>95% CI</i>	3.1–4.9	—	0.5–3.1	9.3–15.0
Females	% of people	7.0	0.0	2.2	19.7
	<i>95% CI</i>	6.0–8.2	—	1.3–3.6	16.7–23.2
Residential location					
Capital city	% of people	4.0	0.0	1.1	12.9
	<i>95% CI</i>	3.3–4.8	—	0.6–1.8	10.6–15.8
Other places	% of people	8.1	0.0	2.9	20.1
	<i>95% CI</i>	6.6–9.8	—	1.4–6.1	16.6–24.3
Postcode socioeconomic status					
Lowest	% of people	8.2	0.0	2.5	21.3
	<i>95% CI</i>	6.8–9.9	—	1.0–5.9	17.7–25.4
Middle	% of people	5.4	0.0	1.9	17.4
	<i>95% CI</i>	4.3–6.8	—	0.9–4.0	14.2–21.1
Highest	% of people	2.9	0.0	0.9	8.7
	<i>95% CI</i>	2.1–3.9	—	0.4–2.0	6.1–12.3
Government health card					
Health care card or pensioner concession card	% of people	15.4	0.0	6.2	24.1
	<i>95% CI</i>	13.3–17.8	—	2.7–13.5	20.9–27.7
Neither card	% of people	2.0	0.0	0.9	8.0
	<i>95% CI</i>	1.6–2.5	—	0.6–1.6	6.2–10.3
Place of last dental visit					
Cardholder/Public	% of people	14.5	0.0	10.9	29.8
	<i>95% CI</i>	10.8–19.3	—	5.0–22.0	22.5–38.4
Cardholder/Non-public	% of people	15.8	0.0	3.2	22.9
	<i>95% CI</i>	13.2–18.7	—	0.8–11.4	19.2–27.0
Non-cardholder/Non-public	% of people	2.0	0.0	0.9	8.0
	<i>95% CI</i>	1.6–2.5	—	0.6–1.6	6.2–10.3
Insured	% of people	2.3	0.0	0.5	7.7
	<i>95% CI</i>	1.7–3.0	—	0.2–1.2	5.7–10.4
Uninsured	% of people	8.4	0.0	3.3	21.9
	<i>95% CI</i>	7.2–9.8	—	1.9–5.7	19.1–25.1

(a) 95% CI = 95% confidence interval for estimated percentage.

Inadequate natural dentition among dentate people

Adults who have approximately 20 teeth or more usually have satisfactory chewing function (Elias & Sheiham 1998), diet and nutritional status (Sheiham et al. 2002), whereas people with fewer teeth are more likely to suffer impaired quality of oral health (McGrath & Bedi 2002). In NSAOH, people were asked during the interview to report either the number of remaining teeth or the number of missing teeth in their upper jaw and lower jaw. Responses were used to classify people as having an inadequate natural dentition if they reported having fewer than 21 natural teeth, the same threshold that has been reported for the UK population. In NSW, 11.7% of dentate adults had fewer than 21 teeth (Table 6), which was almost identical to the national figure of 11.4% (Slade et al. 2007).

Key findings

- The prevalence of an inadequate natural dentition was strongly associated with age, occurring in fewer than 1% of people aged 15–34 years but affecting approximately one-third of dentate people aged 55 years or more.
- Differences in prevalence between males and females were small and statistically non-significant, both for the NSW population as a whole and within the three age groups.
- Similarly, there were no meaningful differences in prevalence between people living in Sydney compared with the remainder of NSW.
- People living in postcodes with low socioeconomic status were approximately twice as likely to report an inadequate natural dentition than people in postcodes of high socioeconomic status. The differences were significant for the whole NSW population and within the two older age groups. People living in postcodes with middle socioeconomic status had prevalence rates that were intermediate between the other two groups.
- Recording the most pronounced differences in prevalence, government health cardholders within the two older age groups were approximately twice as likely as non-cardholders to have an inadequate natural dentition.
- Within the population of government health cardholders, there was a tendency for age-group-specific prevalence to be higher for those whose last dental visit was to the public sector than for those who attended a private dentist. However, the difference was not statistically significant.
- Within the two older age groups and all ages combined, large differences in prevalence were observed between the insured and uninsured.

Discussion

A threshold of fewer than 21 teeth is used here as an indicator of likely impairments in oral function, nutrition and quality of life, rather than a cardinal sign of those problems. As observed for complete tooth loss, there was a pronounced age gradient in prevalence of an inadequate natural dentition. Because of this age association, valid comparisons between other sociodemographic groups should only be made within age groups. Those comparisons reveal that prevalence was associated with postcode socioeconomic status, government health cardholder status and dental insurance. However, unlike the pattern observed for complete tooth loss, prevalence of an inadequate natural dentition did not vary significantly between the sexes or between Sydney and the rest of NSW.

Table 6: Percentage of people with fewer than 21 teeth

		Population: dentate people Age (years)			
		All ages	15–34	35–54	≥55
All people	Per cent of people	11.7	0.2	6.3	34.7
	<i>95% CI^(a)</i>	<i>10.5–13.1</i>	<i>0.1–0.6</i>	<i>4.8–8.2</i>	<i>31.3–38.3</i>
Sex					
Males	% of people	10.7	0.2	6.4	31.1
	<i>95% CI</i>	<i>9.1–12.6</i>	<i>0.0–1.1</i>	<i>4.4–9.4</i>	<i>26.8–35.8</i>
Females	% of people	12.8	0.2	6.2	38.2
	<i>95% CI</i>	<i>11.3–14.4</i>	<i>0.1–0.9</i>	<i>4.5–8.4</i>	<i>34.2–42.4</i>
Residential location					
Capital city	% of people	10.3	0.2	6.0	31.9
	<i>95% CI</i>	<i>8.7–12.1</i>	<i>0.1–0.9</i>	<i>4.3–8.5</i>	<i>27.5–36.8</i>
Other places	% of people	14.3	0.2	6.7	38.7
	<i>95% CI</i>	<i>12.4–16.4</i>	<i>0.0–1.1</i>	<i>4.4–10.1</i>	<i>33.6–44.1</i>
Postcode socioeconomic status					
Lowest	% of people	15.5	0.2	7.3	43.0
	<i>95% CI</i>	<i>13.1–18.2</i>	<i>0.0–1.2</i>	<i>5.0–10.6</i>	<i>37.5–48.6</i>
Middle	% of people	12.2	0.4	7.7	38.5
	<i>95% CI</i>	<i>10.1–14.6</i>	<i>0.1–1.6</i>	<i>5.1–11.5</i>	<i>32.7–44.5</i>
Highest	% of people	7.8	0.0	3.9	23.3
	<i>95% CI</i>	<i>6.4–9.6</i>	—	<i>2.1–7.2</i>	<i>19.5–27.7</i>
Government health card					
Health care card or pensioner concession card	% of people	31.3	0.8	20.1	50.4
	<i>95% CI</i>	<i>27.9–34.8</i>	<i>0.2–3.1</i>	<i>14.3–27.7</i>	<i>45.4–55.4</i>
Neither card	% of people	6.0	0.1	4.0	22.2
	<i>95% CI</i>	<i>5.1–7.1</i>	<i>0.0–0.7</i>	<i>2.9–5.6</i>	<i>18.8–26.1</i>
Place of last dental visit					
Cardholder/Public	% of people	27.1	0.7	31.2	56.4
	<i>95% CI</i>	<i>20.9–34.4</i>	<i>0.1–5.0</i>	<i>18.1–48.2</i>	<i>43.4–68.6</i>
Cardholder/Non-public	% of people	32.9	0.8	13.7	49.2
	<i>95% CI</i>	<i>28.5–37.6</i>	<i>0.1–5.5</i>	<i>8.1–22.3</i>	<i>43.6–54.8</i>
Non-cardholder/Non-public	% of people	6.0	0.1	4.0	22.2
	<i>95% CI</i>	<i>5.1–7.1</i>	<i>0.0–0.7</i>	<i>2.9–5.6</i>	<i>18.8–26.1</i>
Dental insurance					
Insured	% of people	7.3	0.2	2.8	24.0
	<i>95% CI</i>	<i>6.0–8.8</i>	<i>0.0–1.1</i>	<i>1.7–4.6</i>	<i>19.9–28.6</i>
Uninsured	% of people	16.2	0.2	10.7	43.7
	<i>95% CI</i>	<i>14.2–18.4</i>	<i>0.1–1.0</i>	<i>8.0–14.2</i>	<i>39.5–48.0</i>

(a) 95% CI = 95% confidence interval for estimated percentage.

Denture wearing by dentate people

Removable dentures, also called 'false teeth' are worn to replace missing teeth, with the objective to improve function (for example eating), appearance or both. Whereas virtually all edentulous people wear dentures, the decision of dentate people to wear dentures is influenced by numerous factors in addition to the number and location of missing teeth. In NSAOH, removable denture wearing was assessed during the interview by asking two similar questions, 'Do you have a denture or false teeth for your upper (lower) jaw?'. There were 15.9% of dentate adults in NSW who reported wearing one or two dentures (Table 7), a figure that was similar to the 14.9% (Slade et al. 2007) reported nationally.

Key findings

- The frequency of denture wearing was strongly associated with age, ranging from 1.4% among 15–34-year-olds to 43.5% among people aged 55 years or more.
- There were small and statistically non-significant differences between the sexes in the percentage of denture wearers, both within age groups and among all ages combined.
- Similarly, comparisons within age groups between Sydney and the rest of NSW revealed statistically non-significant differences in the percentage of dentate people wearing dentures.
- There was a pronounced socioeconomic gradient in denture wearing, with higher percentages observed among people living in postcodes with low socioeconomic status compared with high socioeconomic status, a pattern that was apparent within each age group and the two older age groups.
- Pronounced differences of a similar magnitude were seen between government health cardholders, one-third of whom wore dentures, and people who did not have a government health card. The difference was statistically significant for all ages combined and for the two older age groups.
- Within the population of government health cardholders, there was inconsistency among age groups in the pattern of difference between people who attended the public sector and those who attended non-public sources of dental care.
- People without dental insurance were more likely to wear dentures than the insured, and the difference was statistically significant for all ages combined and for the two older age groups.

Discussion

The percentage of dentate adults in NSW who wore dentures (15.9%) exceeded the percentage with fewer than 21 natural teeth (11.7%), illustrating that the decision to wear dentures is dictated by factors other than the number of missing teeth. In general, however, sociodemographic variation in frequency of denture wearing was of a similar direction and magnitude to sociodemographic variation in prevalence of an inadequate natural dentition. The exception was residential location outside Sydney, which, in all ages combined, was associated with increased frequency of denture wearing but not associated with prevalence of an inadequate dentition. However, both conditions are strongly associated with age, and more valid comparisons are made within age groups. Those comparisons revealed only small and statistically non-significant differences between Sydney and the rest of NSW for both conditions.

Table 7: Percentage of dentate people who wear denture(s)

		Population: dentate people Age (years)			
		All ages	15–34	35–54	≥55
All people	Per cent of people	15.9	1.4	10.0	43.5
	95% CI ^(a)	14.5–17.4	0.6–3.2	8.2–12.1	40.4–46.7
Sex					
Males	% of people	15.3	1.8	9.9	41.4
	95% CI	13.3–17.5	0.5–5.8	7.3–13.4	37.0–45.9
Females	% of people	16.5	1.0	10.0	45.6
	95% CI	14.7–18.4	0.4–2.5	7.9–12.6	41.3–50.1
Residential location					
Capital city	% of people	14.1	0.9	9.2	41.4
	95% CI	12.5–15.9	0.4–2.2	6.9–12.1	37.4–45.5
Other places	% of people	19.1	2.3	11.3	46.6
	95% CI	16.5–22.0	0.6–8.7	8.7–14.7	41.8–51.5
Postcode socioeconomic status					
Lowest	% of people	21.8	3.2	13.7	52.8
	95% CI	19.3–24.5	1.0–9.6	10.7–17.4	49.0–56.7
Middle	% of people	14.4	0.9	9.1	44.7
	95% CI	12.4–16.6	0.3–2.3	6.3–13.0	39.6–50.0
Highest	% of people	12.0	0.5	7.4	33.4
	95% CI	10.0–14.4	0.1–3.1	4.6–11.6	28.7–38.4
Government health card					
Health care card or pensioner concession card	% of people	33.4	2.9	13.5	56.2
	95% CI	29.6–37.4	1.0–7.5	9.2–19.5	51.9–60.5
Neither card	% of people	10.8	1.1	9.4	33.4
	95% CI	9.4–12.3	0.4–3.5	7.5–11.7	29.3–37.9
Place of last dental visit					
Cardholder/Public	% of people	25.5	5.4	9.9	65.1
	95% CI	20.1–31.7	1.7–16.0	5.1–18.5	54.2–74.6
Cardholder/Non-public	% of people	36.5	0.8	15.6	54.5
	95% CI	31.9–41.4	0.1–5.5	9.7–24.2	49.4–59.4
Non-cardholder/Non-public	% of people	10.8	1.1	9.4	33.4
	95% CI	9.4–12.3	0.4–3.5	7.5–11.7	29.3–37.9
Dental insurance					
Insured	% of people	12.5	0.9	7.6	35.4
	95% CI	10.7–14.5	0.3–2.8	5.6–10.3	30.3–40.8
Uninsured	% of people	19.5	1.9	13.0	50.4
	95% CI	17.2–22.0	0.7–5.2	10.3–16.2	46.5–54.3

(a) 95% CI = 95% confidence interval for estimated percentage.

Average number of teeth per person missing due to pathology

During NSAOH examinations of people aged less than 45 years, dentists counted the number of teeth judged to be missing due to decay or gum disease; for older age groups, dentists counted the number of teeth missing for any reason. The distinction according to age was made because often it is very difficult to judge in older people whether teeth have been extracted because of decay, gum disease or other causes (for example orthodontic reasons), or whether the teeth never developed or remain unerupted. Instead, the convention is to assume that teeth not present among people aged 45 years or more are missing due to pathology.

In NSW, dentate people had an average of 4.9 teeth per person missing due to pathology (Table 8), a figure that was similar to the national average of 4.5% (Slade et al. 2007).

Key findings

- The average number of missing teeth per person was strongly associated with age, ranging from less than 1 among 15–34-year-olds to 11.5 among people aged 55 years or more.
- There was little difference between males and females, and the differences were inconsistent between age groups.
- While there was a tendency for the number of missing teeth to be lower in Sydney than the rest of NSW, the differences were not statistically significant.
- A socioeconomic gradient was observed only in the oldest age group, whereby people living in postcodes with low socioeconomic status had more teeth missing due to pathology, on average, than those living in postcodes with high socioeconomic status.
- Average levels of tooth loss tended to be higher among government health cardholders compared with non-government health cardholders, and the differences were statistically significant in the oldest age group and among all ages combined.
- Among the group of government health cardholders, there tended to be higher average levels of tooth loss for people whose last dental visit was in the public sector compared with the private sector, at least among the two older age groups, although the tendency was not statistically significant.
- Tooth loss due to pathology tended to occur at lower levels for people with dental insurance compared with the uninsured, and the difference was statistically significant in the oldest age group and all ages combined.

Discussion

Consistent with findings from preceding tables describing different aspects of tooth loss, the average number of teeth missing due to pathology was very low among the youngest (15–34 years) age group. Furthermore, because average levels of tooth loss were so strongly associated with age, it is prudent to limit inferences about sociodemographic variation to comparisons within age groups. It follows that the most reliable assessments of sociodemographic differences were observed among the oldest age group. As observed for all other measures of tooth loss, the average number of teeth per person missing due to pathology was associated with postcode socioeconomic status, government health cardholder status and dental insurance.

Table 8: Average number of teeth per person missing due to pathology

		Population: dentate people Age (years)			
		All ages	15–34	35–54	≥55
All people	mean	4.9	0.8	4.1	11.5
	95% CI ^(a)	4.4–5.4	0.5–1.2	3.4–4.8	10.6–12.5
Sex					
Males	mean	4.9	1.0	3.9	11.7
	95% CI	4.2–5.7	0.4–1.7	3.2–4.7	10.3–13.1
Females	mean	4.9	0.6	4.3	11.4
	95% CI	4.3–5.5	0.3–0.9	3.4–5.1	10.2–12.6
Residential location					
Capital city	mean	4.4	0.8	3.6	10.9
	95% CI	3.7–5.0	0.4–1.2	3.0–4.3	9.5–12.3
Other places	mean	5.9	0.8	4.9	12.5
	95% CI	4.9–6.8	0.1–1.5	3.5–6.3	11.2–13.8
Postcode socioeconomic status					
Lowest	mean	5.7	0.7	4.2	13.5
	95% CI	4.5–6.8	<0–1.4	2.9–5.5	12.0–14.9
Middle	mean	5.1	0.7	4.4	12.2
	95% CI	4.3–5.9	0.2–1.2	3.1–5.7	10.8–13.5
Highest	mean	4.1	1.0	3.7	9.3
	95% CI	3.3–4.9	0.4–1.6	2.8–4.5	7.5–11.1
Government health card					
Health care card or pensioner concession card	mean	8.4	0.6	6.0	14.1
	95% CI	7.2–9.7	0.2–0.9	4.2–7.8	12.9–15.3
Neither card	mean	3.7	0.9	3.7	9.3
	95% CI	3.2–4.2	0.4–1.3	3.0–4.4	8.1–10.4
Place of last dental visit					
Cardholder/Public	mean	8.3	0.8	7.6	15.6
	95% CI	6.5–10.1	0.2–1.5	5.2–10.1	13.5–17.7
Cardholder/Non-public	mean	8.5	0.4	4.8	13.7
	95% CI	6.8–10.1	0.0–0.8	2.9–6.8	12.2–15.1
Non-cardholder/Non-public	mean	3.7	0.9	3.7	9.3
	95% CI	3.2–4.2	0.4–1.3	3.0–4.4	8.1–10.4
Dental insurance					
Insured	mean	4.1	0.9	3.4	9.5
	95% CI	3.4–4.8	0.3–1.5	2.8–4.0	7.9–11.1
Uninsured	mean	5.7	0.8	4.7	13.2
	95% CI	5.0–6.5	0.4–1.2	3.7–5.7	12.2–14.3

(a) 95% CI = 95% confidence interval for estimated mean.

Prevalence of untreated coronal decay

The prevalence of untreated coronal dental decay is reported in Table 9 as the percentage of dentate people who have at least one or more decayed surfaces on the crowns of their teeth. Untreated coronal decay reflects both the prevalence of dental decay in the population and access to dental care for treatment.

The prevalence of untreated coronal decay in NSW was 27.1% (Table 9), which is slightly higher than the national estimate of 25.5% (Slade et al. 2007).

Key findings

- The prevalence of untreated coronal decay was not associated with age.
- Among people of all ages, prevalence varied significantly by location, socioeconomic status, government health cardholder status, place of last dental visit and dental insurance.
- The highest prevalence was seen among government health cardholders who last visited a public clinic (49.2%) and the lowest among those living in an area of high socioeconomic status (19.0%).
- Significantly more people living outside Sydney (37.4%) had untreated coronal decay than those in Sydney (21.4%).
- Significantly fewer residents in areas of high socioeconomic status had untreated coronal decay (19.0%) than in those of middle or low socioeconomic status (31.1% and 32.9% respectively).
- Prevalence was significantly higher among government health cardholders (36.9%) compared with non-government health cardholders (23.8%).
- Government health cardholders who last visited a non-public practitioner had the lowest prevalence (23.8%) of untreated coronal decay, significantly lower than that of government health cardholders whose last visit was at a public clinic (49.2%).
- Dental insurance was significantly associated with untreated dental decay, with uninsured people having 1.6 times the prevalence compared with insured people (33.2% versus 20.7%).
- Among people aged 55 years or more, significant differences were found in relation to insurance status.
- Among people aged 35–54 years, significant differences were associated with government health cardholder status, place of last dental visit and insurance status.
- Among people aged 15–34 years, significant differences were found in relation to location.

Discussion

Because untreated decay is associated with poorer access to treatment services, prevalence was associated with social disadvantage and location.

In summary, over one-quarter of all people in NSW had untreated coronal decay, which was more prevalent in groups with higher social disadvantage.

Table 9: Percentage of people with untreated coronal decay

		Population: dentate people Age (years)			
		All ages	15–34	35–54	≥55
All people	Per cent of people	27.1	23.8	28.5	29.8
	<i>95% CI^(a)</i>	23.6–30.9	17.4–31.5	23.8–33.7	25.2–34.8
Sex					
Males	% of people	28.9	20.8	34.7	32.3
	<i>95% CI</i>	24.0–34.4	12.2–33.2	26.8–43.5	25.5–39.9
Females	% of people	25.3	26.7	22.3	27.3
	<i>95% CI</i>	20.9–30.1	18.5–37.0	16.4–29.5	21.2–34.3
Residential location					
Capital city	% of people	21.4	16.1	23.4	26.6
	<i>95% CI</i>	17.6–25.6	9.5–25.8	18.2–29.5	20.2–34.1
Other places	% of people	37.4	39.8	37.6	34.5
	<i>95% CI</i>	30.7–44.6	27.1–54.2	28.9–47.1	28.7–40.8
Postcode socioeconomic status					
Lowest	% of people	32.9	30.5	35.4	32.2
	<i>95% CI</i>	26.6–39.9	20.4–43.0	27.4–44.3	25.0–40.3
Middle	% of people	31.1	30.7	31.0	31.6
	<i>95% CI</i>	24.4–38.6	19.1–45.5	22.4–41.2	23.7–40.6
Highest	% of people	19.0	13.8	19.5	26.1
	<i>95% CI</i>	14.3–24.7	6.1–28.1	13.0–28.3	18.3–35.8
Government health card					
Health care card or pensioner concession card	% of people	36.9	25.8	56.6	33.3
	<i>95% CI</i>	30.0–44.3	13.2–44.1	42.9–69.4	26.9–40.3
Neither card	% of people	23.8	23.6	22.5	26.6
	<i>95% CI</i>	19.8–28.3	16.7–32.3	17.6–28.5	20.4–34.0
Place of last dental visit					
Cardholder/Public	% of people	49.2	43.6	71.2	33.3
	<i>95% CI</i>	37.0–61.5	19.7–70.9	48.0–86.9	22.1–46.8
Cardholder/Non-public	% of people	31.5	16.2	46.1	33.3
	<i>95% CI</i>	23.7–40.4	5.3–40.0	29.5–63.7	25.5–42.0
Non-cardholder/Non-public	% of people	23.8	23.6	22.5	26.6
	<i>95% CI</i>	19.8–28.3	16.7–32.3	17.6–28.5	20.4–34.0
Dental insurance					
Insured	% of people	20.7	24.0	16.6	21.8
	<i>95% CI</i>	15.6–26.8	14.1–37.9	10.9–24.4	15.5–29.8
Uninsured	% of people	33.2	24.8	38.9	36.2
	<i>95% CI</i>	28.6–38.2	17.0–34.5	30.7–47.7	30.5–42.2

(a) 95% CI = 95% confidence interval for estimated percentage.

Percentage of people with untreated root decay

The prevalence of untreated root decay is reported as the percentage of people who had at least one natural tooth and one or more surfaces of the roots of their teeth decayed. Decay of the root surface requires that it be exposed in the mouth, usually by recession of the gums. The prevalence of untreated root decay in NSW was 7.9% (Table 10), which is slightly higher than for the Australian population figure (6.7%) (Slade et al. 2007).

Key findings

- There was a 58.7-fold relative difference between prevalence of untreated root decay in those aged 55 years or more and those aged 15–34 years (17.6% versus 0.3%); and a greater than 2-fold difference between those aged 55 years or more and those in the 35–54 years age group (17.6% versus 8.3%).
- Among people of all ages, the highest prevalence was seen in government health cardholders who last visited a public clinic (23.1%) and the lowest in those who hold private dental insurance (4.3%).
- Significantly fewer residents in areas of high socioeconomic status had untreated root decay (4.8%) than in those of low socioeconomic status (10.3%).
- Prevalence was significantly higher among government health cardholders (14.2%) compared with non-government health cardholders (5.7%).
- Non-government health cardholders who last visited a non-public practitioner recorded the lowest prevalence (5.7%) of untreated root decay. This was significantly lower than that of government health cardholders whose last visit was at a public clinic (23.1%).
- Uninsured people had 2.6 times the prevalence of untreated root decay compared with insured people (11.2% versus 4.3%).

Discussion

The association of root decay with gum recession more commonly seen in older people explains the strong relationship of untreated root decay with age. Untreated disease reflects access to timely dental care and therefore untreated root decay is more common among people who are socially disadvantaged.

In summary, prevalence of root decay varied by age, socioeconomic status, government health cardholder status, place of last dental visit and insurance status.

Table 10: Percentage of people with untreated root decay

		Population: dentate people			
		Age (years)			
		All ages	15–34	35–54	≥55
All people	Per cent of people	7.9	0.3	8.3	17.6
	<i>95% CI^(a)</i>	6.5–9.6	0.0–2.0	5.9–11.7	13.7–22.3
Sex					
Males	% of people	8.9	0.0	9.4	20.3
	<i>95% CI</i>	6.6–11.8	—	5.8–15.0	15.0–27.0
Females	% of people	6.9	0.6	7.2	14.9
	<i>95% CI</i>	5.0–9.5	0.1–4.0	4.2–12.3	10.0–21.6
Residential location					
Capital city	% of people	6.5	0.4	6.5	16.1
	<i>95% CI</i>	5.0–8.5	0.1–2.9	3.7–10.9	11.5–22.0
Other places	% of people	10.3	0.0	11.7	19.8
	<i>95% CI</i>	7.7–13.5	—	7.6–17.4	13.6–28.0
Postcode socioeconomic status					
Lowest	% of people	10.3	1.0	11.9	18.9
	<i>95% CI</i>	7.6–13.9	0.2–6.7	7.5–18.5	12.4–27.8
Middle	% of people	9.1	0.0	7.9	23.1
	<i>95% CI</i>	6.7–12.3	—	4.2–14.5	16.4–31.5
Highest	% of people	4.8	0.0	5.3	11.6
	<i>95% CI</i>	3.2–7.2	—	2.5–10.9	6.8–19.1
Government health card					
Health care card or pensioner concession card	% of people	14.2	0.0	19.8	19.4
	<i>95% CI</i>	10.5–18.9	—	10.1–35.3	14.9–24.9
Neither card	% of people	5.7	0.4	5.9	16.0
	<i>95% CI</i>	4.2–7.6	0.1–2.5	3.7–9.2	10.9–22.7
Place of last dental visit					
Cardholder/Public	% of people	23.1	0.0	43.9	24.1
	<i>95% CI</i>	14.2–35.2	—	23.3–66.7	14.7–36.9
Cardholder/Non-public	% of people	10.3	0.0	2.6	18.1
	<i>95% CI</i>	7.0–14.9	—	1.0–6.3	12.6–25.3
Non-cardholder/Non-public	% of people	5.7	0.4	5.9	16.0
	<i>95% CI</i>	4.2–7.6	0.1–2.5	3.7–9.2	10.9–22.7
Dental insurance					
Insured	% of people	4.3	0.0	3.0	12.1
	<i>95% CI</i>	2.7–6.6	—	1.1–7.4	7.7–18.4
Uninsured	% of people	11.2	0.6	13.0	22.1
	<i>95% CI</i>	9.1–13.8	0.1–3.9	8.9–18.7	16.8–28.5

(a) 95% CI = 95% confidence interval for estimated percentage.

Percentage of people with one or more filled teeth

Fillings for treatment of tooth decay leave permanent marks on the teeth and are one measure of a person's lifetime experience of decay. Filled teeth also indicate patterns of dental treatment and access to dental care. The prevalence of filled teeth in NSW was 79.6% (Table 11), which is slightly higher than for the overall Australian population figure (83.9%) (Slade et al. 2007).

Key findings

- Prevalence of filled teeth was significantly associated with age; among those aged 55 years or more and those aged 35–54 years, it was over 1.5 times that of those in the 15–34 years age group (91.2% and 91.5% versus 59.2% respectively).
- Among people of all ages, the highest prevalence was seen among those with private dental insurance (86.8%) and the lowest among government health cardholders who last attended a public clinic (72.9%).
- Dental insurance was significantly associated with filled teeth, with insured people having 1.2 times the prevalence compared with uninsured people (86.8% versus 74.2%).

Discussion

The percentage of people with filled teeth relates to lifetime experience of dental decay, and hence is associated with age. Prevalence also reflects access to timely dental care, and type of care used to treat caries being a restoration rather than an extraction; hence, there is an association with private insurance which facilitates access to dental care.

In summary, the percentage of people with filled teeth was associated with age and dental insurance.

Table 11: Percentage of people with one or more filled teeth

		Population: dentate people Age (years)			
		All ages	15–34	35–54	≥55
All people	Per cent of people	79.6	59.2	91.5	91.2
	<i>95% CI^(a)</i>	<i>75.3–83.3</i>	<i>50.1–67.7</i>	<i>87.4–94.4</i>	<i>87.5–93.9</i>
Sex					
Males	% of people	79.0	61.1	89.1	90.0
	<i>95% CI</i>	<i>71.0–85.2</i>	<i>45.0–75.0</i>	<i>82.1–93.6</i>	<i>84.3–93.9</i>
Females	% of people	80.2	57.2	93.9	92.3
	<i>95% CI</i>	<i>76.1–83.8</i>	<i>47.9–66.1</i>	<i>90.0–96.4</i>	<i>86.0–95.9</i>
Residential location					
Capital city	% of people	81.2	64.9	90.7	92.3
	<i>95% CI</i>	<i>75.8–85.5</i>	<i>53.8–74.6</i>	<i>84.9–94.4</i>	<i>86.4–95.8</i>
Other places	% of people	76.8	47.3	92.9	89.5
	<i>95% CI</i>	<i>69.1–83.1</i>	<i>32.4–62.6</i>	<i>86.5–96.4</i>	<i>84.8–92.9</i>
Postcode socioeconomic status					
Lowest	% of people	75.2	41.1	92.5	91.0
	<i>95% CI</i>	<i>65.7–82.8</i>	<i>25.4–58.9</i>	<i>85.9–96.2</i>	<i>85.1–94.6</i>
Middle	% of people	82.8	70.5	91.0	87.9
	<i>95% CI</i>	<i>76.6–87.6</i>	<i>55.6–82.1</i>	<i>82.1–95.8</i>	<i>79.9–93.0</i>
Highest	% of people	80.4	62.4	91.0	94.3
	<i>95% CI</i>	<i>73.3–86.0</i>	<i>49.7–73.5</i>	<i>83.5–95.3</i>	<i>86.6–97.7</i>
Government health card					
Health care card or pensioner concession card	% of people	78.0	50.6	93.4	85.8
	<i>95% CI</i>	<i>70.9–83.7</i>	<i>32.7–68.4</i>	<i>78.6–98.2</i>	<i>78.5–90.8</i>
Neither card	% of people	80.7	62.3	91.1	96.0
	<i>95% CI</i>	<i>75.0–85.3</i>	<i>51.2–72.2</i>	<i>86.3–94.3</i>	<i>92.3–98.0</i>
Place of last dental visit					
Cardholder/Public	% of people	72.9	35.7	94.4	85.8
	<i>95% CI</i>	<i>59.5–83.1</i>	<i>14.9–63.7</i>	<i>83.2–98.3</i>	<i>70.2–93.9</i>
Cardholder/Non-public	% of people	80.2	58.5	92.8	85.7
	<i>95% CI</i>	<i>71.0–87.0</i>	<i>34.1–79.4</i>	<i>63.4–99.0</i>	<i>76.9–91.6</i>
Non-cardholder/Non-public	% of people	80.7	62.3	91.1	96.0
	<i>95% CI</i>	<i>75.0–85.3</i>	<i>51.2–72.2</i>	<i>86.3–94.3</i>	<i>92.3–98.0</i>
Dental insurance					
Insured	% of people	86.8	73.8	93.5	95.6
	<i>95% CI</i>	<i>81.2–91.0</i>	<i>60.5–83.8</i>	<i>88.2–96.5</i>	<i>89.5–98.2</i>
Uninsured	% of people	74.2	46.8	89.8	87.6
	<i>95% CI</i>	<i>67.5–79.9</i>	<i>34.4–59.7</i>	<i>82.8–94.2</i>	<i>81.8–91.7</i>

(a) 95% CI = 95% confidence interval for estimated percentage.

Average number of decayed, missing and filled teeth per person

The number of decayed, missing because of pathology, and filled teeth (DMFT) reflects a person's lifetime experience of dental caries. In this survey all missing teeth in people aged 45 years or more were counted as missing due to pathology, while for people aged less than 45 years, the count only included teeth where the examiner judged that dental decay or gum disease was the likely reason for the extraction.

The average DMFT number in NSW was 12.8 (Table 12), which is the same as that for the overall Australian population (12.8 teeth) (Slade et al. 2007).

Key findings

- The average number of affected teeth was significantly associated with age, being highest in people aged 55 years or more (22.7 teeth). This was 1.6 times that of those aged 35–44 years (13.9 teeth) and five times that of the 15–34 years age group (4.4 teeth).
- Among people of all ages, the highest average was seen in government health cardholders who last attended a non-public clinic (16.8 teeth) and the lowest among non-government health cardholders who last visited a non-public clinic (11.5 teeth).
- Government health cardholder status was associated with average number of teeth with caries experience, with cardholders having 1.4 times the prevalence compared with non-cardholders (16.6 versus 11.5 teeth).
- The average DMFT was significantly associated with place of last dental visit, with government health cardholders who last visited a public clinic and those who visited a non-public practitioner having significantly higher scores than non-government health cardholders whose last visit was at a private clinic (16.2 and 16.8 versus 11.5 teeth respectively).
- Among people aged 55 years or more, a significant difference in average decay experience was found in relation to government health cardholder status.
- Among people aged 35–54 years, a significant difference in average DMFT score was found in relation to government health cardholder status.

Discussion

The average number of teeth with caries experience over a lifetime is a cumulative score, and hence is strongly associated with age. In the oldest age group (55 years or more), most of a person's 32 teeth had experienced dental caries. Disease experience is related to disadvantage, as evidenced by associations with government health cardholder status and place of last dental visit.

In summary, average DMFT scores were strongly related to age, but were also associated with some measures of disadvantage.

Table 12: Average number of decayed, missing or filled teeth per person

		Population: dentate people Age (years)			
		All ages	15–34	35–54	≥55
All people	mean	12.8	4.4	13.9	22.7
	95% CI ^(a)	11.9–13.7	3.4–5.4	13.0–14.8	22.2–23.2
Sex					
Males	mean	12.5	4.9	13.5	21.9
	95% CI	11.1–14.0	3.0–6.7	12.2–14.8	21.1–22.7
Females	mean	13.0	3.8	14.4	23.5
	95% CI	11.9–14.2	3.1–4.6	13.2–15.5	22.8–24.2
Residential location					
Capital city	mean	12.2	4.2	13.4	22.5
	95% CI	11.1–13.3	3.2–5.2	12.4–14.5	21.8–23.2
Other places	mean	13.9	4.7	14.8	22.9
	95% CI	12.3–15.5	2.4–7.0	13.1–16.5	22.3–23.6
Postcode socioeconomic status					
Lowest	mean	13.4	4.2	14.0	23.5
	95% CI	11.4–15.5	1.5–6.9	12.3–15.7	22.7–24.3
Middle	mean	12.7	3.8	14.1	22.6
	95% CI	11.4–14.0	2.8–4.8	12.4–15.8	21.9–23.4
Highest	mean	12.4	4.9	13.7	22.0
	95% CI	10.9–13.8	3.5–6.4	12.4–15.0	21.2–22.9
Government health card					
Health care card or pensioner concession card	mean	16.6	4.0	16.7	23.7
	95% CI	14.9–18.3	1.9–6.0	14.9–18.6	23.0–24.4
Neither card	mean	11.5	4.5	13.3	21.8
	95% CI	10.6–12.5	3.4–5.7	12.3–14.4	21.1–22.5
Place of last dental visit					
Cardholder/Public	mean	16.2	4.3	19.3	23.9
	95% CI	13.5–18.9	1.3–7.3	16.1–22.4	22.4–25.4
Cardholder/Non-public	mean	16.8	3.8	14.9	23.6
	95% CI	14.7–18.9	1.1–6.5	13.2–16.6	22.8–24.4
Non-cardholder/Non-public	mean	11.5	4.5	13.3	21.8
	95% CI	10.6–12.5	3.4–5.7	12.3–14.4	21.1–22.5
Dental insurance					
Insured	mean	12.8	5.1	13.3	22.8
	95% CI	11.5–14.1	3.3–6.9	12.2–14.5	22.1–23.5
Uninsured	mean	13.1	3.9	14.4	22.7
	95% CI	11.9–14.2	2.8–5.0	13.2–15.7	21.9–23.4

(a) 95% CI = 95% confidence interval for estimated mean.

Prevalence of moderate or severe periodontitis

A case definition of periodontitis has been developed jointly by the US Centers for Disease Control and Prevention (CDC) and the American Academy of Periodontology (AAP) to describe prevalence of moderate and severe periodontitis. The CDC–AAP defines moderate periodontitis as the presence of either two sites between adjacent teeth where the gum has lost its attachment to the tooth for 4 mm or more, or at least two such sites that have pockets of 5 mm or more. Severe periodontitis has been defined as having at least two sites between adjacent teeth where the gum has lost its attachment to the tooth for 6 mm or more, and there is at least one pocket of 5 mm or greater depth. Table 13 reports estimates of combined moderate and severe periodontitis.

In NSW, a total of 23.0% of the dentate population had moderate or severe periodontitis (Table 13), which was similar to the national estimate of 22.9% (Slade et al. 2007).

Key findings

- The prevalence of moderate or severe periodontitis was strongly associated with age in NSW, being 6% in 15–34-year-old adults but affecting 44% of those aged 55 years or more.
- Government health cardholders were almost twice as likely to have periodontitis, both for all ages combined and for people in the 35–54 years age group. This difference remained even after stratification by place of the last dental visit.
- People with no private dental insurance were significantly more likely to have periodontitis. The difference was also significant for those aged 55 years or more.

Discussion

Components of periodontal disease measurement reflect both concurrent disease state and historical accumulation of the disease. Therefore, a strong association with age was fully expected. Because periodontitis was more prevalent in the middle-aged and older segments of the population, comparisons between population groups were observed most clearly in those age groups.

In summary, moderate or severe periodontitis affected one-quarter of the NSW adult population, with the highest proportion of those affected being in the oldest age group. The disease was most likely to be observed in the socioeconomically disadvantaged groups.

Table 13: Percentage of people with moderate or severe periodontitis

		Population: dentate people			
		Age (years)			
		All ages	15–34	35–54	≥55
All people	Per cent of people	23.0	6.3	26.1	44.1
	<i>95% CI^(a)</i>	<i>19.7–26.7</i>	<i>3.0–12.6</i>	<i>21.1–31.9</i>	<i>38.6–49.8</i>
Sex					
Males	% of people	27.1	8.8	32.3	49.3
	<i>95% CI</i>	<i>21.5–33.5</i>	<i>3.4–20.9</i>	<i>24.7–41.0</i>	<i>40.5–58.2</i>
Females	% of people	18.8	3.5	19.7	39.3
	<i>95% CI</i>	<i>15.4–22.7</i>	<i>1.6–7.6</i>	<i>14.4–26.4</i>	<i>32.2–46.9</i>
Residential location					
Capital city	% of people	25.2	6.3	30.1	51.3
	<i>95% CI</i>	<i>20.7–30.2</i>	<i>2.5–15.0</i>	<i>23.2–37.9</i>	<i>43.5–59.1</i>
Other places	% of people	18.9	6.3	19.1	33.6
	<i>95% CI</i>	<i>14.5–24.2</i>	<i>1.9–18.3</i>	<i>13.0–27.2</i>	<i>26.5–41.5</i>
Postcode socioeconomic status					
Lowest	% of people	19.8	1.2	22.9	37.1
	<i>95% CI</i>	<i>14.6–26.3</i>	<i>0.3–5.0</i>	<i>14.3–34.6</i>	<i>28.3–46.9</i>
Middle	% of people	25.4	7.7	29.4	47.8
	<i>95% CI</i>	<i>20.1–31.5</i>	<i>2.8–19.3</i>	<i>22.0–38.1</i>	<i>39.4–56.3</i>
Highest	% of people	23.4	8.1	26.0	47.2
	<i>95% CI</i>	<i>17.6–30.3</i>	<i>2.6–22.4</i>	<i>17.3–37.1</i>	<i>36.4–58.4</i>
Government health card					
Health care card or pensioner concession card	% of people	36.2	13.6	42.6	47.6
	<i>95% CI</i>	<i>28.4–44.6</i>	<i>4.3–35.6</i>	<i>28.9–57.5</i>	<i>38.5–56.8</i>
Neither card	% of people	18.7	4.6	22.7	41.1
	<i>95% CI</i>	<i>15.8–21.9</i>	<i>1.9–10.5</i>	<i>18.3–27.9</i>	<i>34.0–48.7</i>
Place of last dental visit					
Cardholder/Public	% of people	48.1	13.6	64.5	62.4
	<i>95% CI</i>	<i>34.1–62.4</i>	<i>1.9–56.0</i>	<i>41.4–82.3</i>	<i>42.8–78.7</i>
Cardholder/Non-public	% of people	30.9	13.6	26.4	43.2
	<i>95% CI</i>	<i>22.9–40.3</i>	<i>3.2–42.5</i>	<i>13.5–45.2</i>	<i>33.7–53.2</i>
Non-cardholder/Non-public	% of people	18.7	4.6	22.7	41.1
	<i>95% CI</i>	<i>15.8–21.9</i>	<i>1.9–10.5</i>	<i>18.3–27.9</i>	<i>34.0–48.7</i>
Dental insurance					
Insured	% of people	18.1	4.3	21.4	34.9
	<i>95% CI</i>	<i>14.4–22.5</i>	<i>1.3–13.6</i>	<i>15.5–28.8</i>	<i>28.0–42.5</i>
Uninsured	% of people	27.9	8.6	30.3	51.7
	<i>95% CI</i>	<i>23.1–33.3</i>	<i>3.5–19.5</i>	<i>23.4–38.1</i>	<i>43.4–60.0</i>

(a) 95% CI = 95% confidence interval for estimated percentage.

Prevalence of deep pocket depth

Deep periodontal pockets have been defined as 4 mm or more. The depth of the pocket, measured in millimetres using a periodontal probe, is an indication of the severity of the destructive process.

In NSW, a total of 21.4% of the dentate adult population had at least one site with periodontal pocket depth of 4 mm or more (Table 14), which was higher, but not significantly, than the national estimate of 19.8% (Slade et al. 2007).

Key findings

- The prevalence of deep periodontal pocket varied with age. The 35–54-year-old age group had the highest prevalence (27.5%), which was more than twice as high as the 15–34-year-old group. The oldest age group had an intermediate prevalence compared with the other two groups.
- People aged 55 years or more who reside in the capital city were more likely to have one or more sites with periodontal pocket depth of 4 mm or more.
- There were no statistically significant differences between the other population groups.

Discussion

The depth of periodontal pockets reflects a more current activity of periodontal inflammation. This activity may be more dependent on oral hygiene status, which was found to not vary widely between groups.

In summary, deep periodontal pocketing affected one-fifth of the NSW dentate population. The middle-aged group (35–54 years) was the most affected.

Table 14: Percentage of people with 4+ mm periodontal pocket depth

		Population: dentate people Age (years)			
		All ages	15–34	35–54	≥55
All people	Per cent of people	21.4	13.0	27.5	24.9
	95% CI ^(a)	17.8–25.5	7.7–21.2	21.9–33.8	19.7–31.0
Sex					
Males	% of people	24.8	16.1	33.8	24.6
	95% CI	18.7–32.1	7.6–30.7	24.9–44.1	17.9–32.8
Females	% of people	17.9	9.7	20.9	25.2
	95% CI	14.6–21.8	4.8–18.6	15.7–27.3	18.5–33.4
Residential location					
Capital city	% of people	25.1	15.0	32.6	30.9
	95% CI	20.2–30.7	8.1–26.1	25.4–40.7	23.0–40.3
Other places	% of people	14.5	8.6	18.3	16.1
	95% CI	9.8–21.0	3.0–22.5	11.5–28.0	11.3–22.4
Postcode socioeconomic status					
Lowest	% of people	14.3	8.2	17.5	16.6
	95% CI	9.6–20.8	2.3–25.5	10.5–27.9	10.9–24.5
Middle	% of people	23.3	12.2	34.1	23.6
	95% CI	17.2–30.6	5.1–26.5	24.1–45.7	15.6–34.1
Highest	% of people	25.3	16.5	30.3	33.6
	95% CI	18.7–33.2	7.5–32.6	21.3–41.1	23.0–46.2
Government health card					
Health care card or pensioner concession card	% of people	23.0	19.9	29.1	21.6
	95% CI	16.4–31.2	8.0–41.6	17.2–44.8	13.4–32.9
Neither card	% of people	21.0	11.5	27.1	27.8
	95% CI	17.0–25.5	6.0–20.9	21.5–33.6	22.3–34.1
Place of last dental visit					
Cardholder/Public	% of people	35.5	28.2	36.2	41.3
	95% CI	23.4–49.7	7.7–65.1	16.6–61.7	23.7–61.5
Cardholder/Non-public	% of people	17.6	16.1	23.9	15.7
	95% CI	11.3–26.2	4.6–43.6	11.5–43.2	8.6–26.9
Non-cardholder/Non-public	% of people	21.0	11.5	27.1	27.8
	95% CI	17.0–25.5	6.0–20.9	21.5–33.6	22.3–34.1
Dental insurance					
Insured	% of people	21.7	10.8	32.3	22.1
	95% CI	16.8–27.6	4.1–25.6	25.3–40.1	16.6–28.7
Uninsured	% of people	21.6	16.0	23.3	27.1
	95% CI	17.1–27.0	8.7–27.6	16.9–31.2	19.3–36.5

(a) 95% CI = 95% confidence interval for estimated percentage.

Prevalence of 4+ mm clinical attachment loss

Clinical attachment loss (CAL) is the loss of supporting periodontal structure around the tooth. Attachment may be lost through gum recession or the development of periodontal pockets from the inflammatory disease periodontitis. In NSAOH, CAL was measured using a combination of gum recession and periodontal probing depth on three sites per tooth.

In NSW, a total of 45% of dentate adults had at least one site with 4 mm or more CAL (Table 15), which was higher, but not significantly, than the national estimate of 42.5% (Slade et al. 2007).

Key findings

- The prevalence of 4+ mm CAL was strongly associated with age in NSW, being 18.3% in 15–34-year-old adults but affecting 76.5% of those aged 55 years or more. Over half of the middle-aged adults also had at least one site with 4+ mm CAL.
- There was a tendency for males to have higher prevalence of CAL of 4+ mm compared with females. This tendency was statistically significant in the older age group.
- Government health cardholders were significantly more likely to have CAL of 4 mm or more compared with people who did not have a government health card.
- People who had their last visit to a public sector facility had significantly higher prevalence of CAL of 4+ mm. The difference was evident for all age groups combined and for the middle-aged group.

Discussion

Clinical attachment loss reflects an accumulation of activity of periodontal inflammation as well as a physiological process in the gums. Therefore, a strong age effect was observed. This condition was almost universal in certain groups of the oldest population. This strong age effect might confound the effect of other attributes depending on data structures within and between groups, such as by residential location.

In summary, clinical attachment loss was highly prevalent in this population. It was more likely to occur in the older population, males and people with lower socioeconomic status.

Table 15: Percentage of people with 4+ mm clinical attachment loss

		Population: dentate people			
		Age (years)			
		All ages	15–34	35–54	≥55
All people	Per cent of people	45.2	18.3	52.0	76.5
	95% CI ^(a)	40.1–50.4	11.8–27.4	46.2–57.8	71.2–81.1
Sex					
Males	% of people	50.6	25.0	56.8	83.8
	95% CI	41.5–59.7	13.5–41.8	47.5–65.6	76.8–89.0
Females	% of people	39.6	11.0	47.1	69.7
	95% CI	34.3–45.2	6.0–19.4	39.6–54.7	62.1–76.5
Residential location					
Capital city	% of people	44.3	19.1	52.0	77.4
	95% CI	38.0–50.8	11.2–30.7	44.4–59.4	69.6–83.7
Other places	% of people	46.7	16.5	52.0	75.1
	95% CI	38.4–55.3	7.1–33.6	43.1–60.8	68.0–81.1
Postcode socioeconomic status					
Lowest	% of people	45.4	12.0	50.8	76.7
	95% CI	36.4–54.6	4.3–29.4	41.1–60.4	68.4–83.4
Middle	% of people	45.5	15.8	57.3	75.1
	95% CI	37.3–54.0	6.9–32.0	46.5–67.4	66.2–82.3
Highest	% of people	44.7	24.1	48.1	77.5
	95% CI	36.0–53.8	12.8–40.6	39.2–57.2	66.3–85.7
Government health card					
Health care card or pensioner concession card	% of people	57.5	16.4	63.1	81.9
	95% CI	48.2–66.3	5.9–37.8	48.2–75.9	75.6–86.8
Neither card	% of people	41.3	19.1	49.7	71.8
	95% CI	35.9–47.0	11.7–29.6	43.6–55.8	63.7–78.8
Place of last dental visit					
Cardholder/Public	% of people	64.5	13.6	84.1	90.7
	95% CI	49.1–77.4	1.9–56.0	62.3–94.4	78.6–96.2
Cardholder/Non-public	% of people	54.4	17.7	47.7	79.3
	95% CI	43.6–64.9	5.2–45.6	31.5–64.3	71.6–85.3
Non-cardholder/Non-public	% of people	41.3	19.1	49.7	71.8
	95% CI	35.9–47.0	11.7–29.6	43.6–55.8	63.7–78.8
Dental insurance					
Insured	% of people	44.8	19.5	52.9	72.4
	95% CI	37.6–52.3	9.7–35.3	45.9–59.8	63.9–79.6
Uninsured	% of people	45.9	16.4	51.2	79.8
	95% CI	39.9–52.0	8.8–28.7	43.5–58.9	72.3–85.6

(a) 95% CI = 95% confidence interval for estimated percentage.

Prevalence of gingival inflammation

The gingival index is a measure of gingivitis, or inflammation of the gums. Gingivitis occurs as a response to the bacteria in plaque accumulation near the gum line. In NSAOH gingivitis was assessed on six index teeth. A gingival index score of 2 or more indicated bleeding on probing or spontaneous bleeding, and was classified as indicating gingival inflammation (gingivitis).

In NSW, a total of 17.2% of the dentate adult population had at least one site with a gingival score of 2 or more (Table 16), which was lower, but not significantly, than the national estimate of 19.7% (Slade et al. 2007).

Key findings

- The prevalence of gingival inflammation was not statistically significant between population groups.
- There was a tendency that people with lower socioeconomic status were more likely to have gingival inflammation. However, the differences were not statistically significant.

Discussion

Gingival inflammation is a condition observed in people of all ages at a similar rate. There was a tendency that people with lower socioeconomic status had higher prevalence of gingival inflammation. However, the differences were small and relatively low numbers of people in each population group made the confidence interval wide, overlapping between groups.

Table 16: Percentage of people with gingival inflammation

		Population: dentate people Age (years)			
		All ages	15–34	35–54	≥55
All people	Per cent of people	17.2	16.3	17.0	19.2
	<i>95% CI^(a)</i>	<i>14.1–20.9</i>	<i>10.6–24.1</i>	<i>13.1–21.7</i>	<i>14.8–24.6</i>
Sex					
Males	% of people	16.5	11.6	19.6	20.0
	<i>95% CI</i>	<i>12.4–21.6</i>	<i>5.2–23.8</i>	<i>13.3–27.7</i>	<i>13.7–28.2</i>
Females	% of people	18.0	21.3	14.3	18.5
	<i>95% CI</i>	<i>13.9–22.9</i>	<i>13.3–32.5</i>	<i>9.6–20.7</i>	<i>13.3–25.2</i>
Residential location					
Capital city	% of people	16.5	16.8	15.5	17.8
	<i>95% CI</i>	<i>12.7–21.3</i>	<i>9.8–27.3</i>	<i>11.0–21.3</i>	<i>12.2–25.2</i>
Other places	% of people	18.6	15.0	19.7	21.3
	<i>95% CI</i>	<i>13.8–24.5</i>	<i>7.8–27.2</i>	<i>13.1–28.4</i>	<i>14.6–29.8</i>
Postcode socioeconomic status					
Lowest	% of people	16.7	12.2	17.5	21.1
	<i>95% CI</i>	<i>12.2–22.5</i>	<i>5.0–26.9</i>	<i>10.7–27.5</i>	<i>13.8–30.7</i>
Middle	% of people	22.8	25.6	18.5	25.1
	<i>95% CI</i>	<i>16.7–30.2</i>	<i>15.6–39.2</i>	<i>12.0–27.5</i>	<i>16.5–36.2</i>
Highest	% of people	12.8	11.4	14.9	12.3
	<i>95% CI</i>	<i>8.8–18.3</i>	<i>4.5–25.8</i>	<i>9.5–22.6</i>	<i>7.6–19.4</i>
Government health card					
Health care card or pensioner concession card	% of people	23.1	17.6	27.0	24.5
	<i>95% CI</i>	<i>16.9–30.7</i>	<i>7.9–34.7</i>	<i>15.2–43.3</i>	<i>16.9–34.1</i>
Neither card	% of people	15.4	16.2	14.9	14.8
	<i>95% CI</i>	<i>11.8–19.9</i>	<i>9.8–25.6</i>	<i>10.9–20.0</i>	<i>10.2–21.0</i>
Place of last dental visit					
Cardholder/Public	% of people	23.3	23.8	28.7	16.8
	<i>95% CI</i>	<i>12.5–39.2</i>	<i>6.0–60.3</i>	<i>11.6–55.2</i>	<i>8.5–30.6</i>
Cardholder/Non-public	% of people	23.0	14.8	25.8	26.7
	<i>95% CI</i>	<i>16.3–31.3</i>	<i>5.6–33.8</i>	<i>13.0–44.9</i>	<i>17.8–37.9</i>
Non-cardholder/Non-public	% of people	15.4	16.2	14.9	14.8
	<i>95% CI</i>	<i>11.8–19.9</i>	<i>9.8–25.6</i>	<i>10.9–20.0</i>	<i>10.2–21.0</i>
Dental insurance					
Insured	% of people	14.3	17.0	11.7	14.3
	<i>95% CI</i>	<i>9.7–20.6</i>	<i>8.8–30.4</i>	<i>7.1–18.6</i>	<i>8.8–22.3</i>
Uninsured	% of people	19.7	15.1	21.6	23.3
	<i>95% CI</i>	<i>15.8–24.3</i>	<i>8.8–24.8</i>	<i>15.8–28.8</i>	<i>16.4–32.0</i>

(a) 95% CI = 95% confidence interval for estimated percentage.

4 Oral health care

Dental attendance within the preceding 12 months

Time since last visiting a dentist is a key indicator of access to dental care. In NSAOH, the time since last dental visit was assessed in the interview by asking 'How long ago did you last see a dental professional about your teeth, dentures or gums?'. Five responses were possible including 'Less than 12 months'. In NSW, 6 out of 10 (60.3%) people aged 15 years or more had visited a dentist within the last 12 months (Table 17). This estimate was not significantly different from the national estimate of 59.4% (Slade et al. 2007).

Key findings

- There was some variation in the percentage of adults visiting within the last 12 months across the three age groups, with adults aged 35–54 years reporting the highest percentage (64.5%).
- A similar percentage of males and females reported visiting a dentist within the last 12 months (57.9% versus 62.6%). Differences by gender were largest in the 15–34 years age group, where 52.0% of males recently visited compared with 62.1% of females.
- People living in Sydney were more likely to report recently visiting a dentist than those living in the rest of NSW (63.3% versus 55.0%). This difference was mainly attributable to those aged 55 years or more where the figures were 65.2% and 50.6%, respectively.
- Across all ages, 68.7% of people living in postcodes with high socioeconomic status had visited a dentist within the last 12 months compared with 53.9% of those living in low socioeconomic postcodes. Within age groups, differences were particularly evident in people aged 55 years or more (73.7% versus 50.0%).
- Government health cardholders were less likely than non-government health cardholders to have attended in the preceding 12 months (51.4% versus 63.5%). Less frequent dental visiting by health cardholders was evident in all age groups, with significant differences for those aged 55 years or more (49.7% versus 68.1%).
- Among government health cardholders, those who attended a public practice at their last visit were almost as likely to have visited a dentist within the last 12 months as people who attended a private practice (48.0% versus 52.7%).
- Insured people were much more likely to have recently visited than uninsured people (74.2% versus 48.3%). Significant differences between insured and uninsured were evident in all age groups.

Discussion

Six out of 10 NSW residents visited a dentist within the last 12 months. Being insured, not having a government health card, residing in the metropolitan region and living in areas of high socioeconomic status were all associated with recent dental visiting. Differences in visiting behaviour were most evident for people aged 55 years or more, with large differences observed by both socioeconomic and insurance status.

Table 17: Percentage of people visiting dentist within last 12 months

		Population: all people Age (years)			
		All ages	15–34	35–54	≥55
All people	Per cent of people	60.3	57.0	64.5	59.0
	95% CI ^(a)	58.3–62.2	53.2–60.8	61.8–67.2	55.5–62.4
Sex					
Males	% of people	57.9	52.0	63.1	58.7
	95% CI	54.9–60.8	45.9–58.1	58.5–67.4	53.8–63.3
Females	% of people	62.6	62.1	66.0	59.3
	95% CI	60.0–65.0	56.9–67.0	62.4–69.4	55.4–63.1
Residential location					
Capital city	% of people	63.3	58.5	66.9	65.2
	95% CI	60.9–65.7	54.2–62.6	63.4–70.3	60.7–69.5
Other places	% of people	55.0	54.0	60.4	50.6
	95% CI	51.7–58.2	46.2–61.5	56.0–64.6	45.0–56.1
Postcode socioeconomic status					
Lowest	% of people	53.9	51.2	60.3	50.0
	95% CI	50.5–57.3	44.4–58.0	55.4–65.1	44.3–55.7
Middle	% of people	58.0	57.6	61.2	54.3
	95% CI	55.0–60.9	50.1–64.7	56.2–66.0	48.5–59.9
Highest	% of people	68.7	61.2	71.9	73.7
	95% CI	66.2–71.1	56.6–65.7	68.0–75.5	69.4–77.6
Government health card					
Health care card or pensioner concession card	% of people	51.4	51.3	56.2	49.7
	95% CI	47.9–54.8	40.4–62.2	49.1–63.0	45.7–53.7
Neither card	% of people	63.5	58.5	66.0	68.1
	95% CI	61.3–65.7	54.3–62.7	63.1–68.9	63.7–72.2
Place of last dental visit					
Cardholder/Public	% of people	48.0	51.0	50.3	43.6
	95% CI	40.2–55.9	35.9–65.9	37.8–62.7	34.0–53.7
Cardholder/Non-public	% of people	52.7	51.6	59.9	51.1
	95% CI	49.0–56.3	38.5–64.4	49.5–69.4	46.7–55.4
Non-cardholder/Non-public	% of people	63.5	58.5	66.0	68.1
	95% CI	61.3–65.7	54.3–62.7	63.1–68.9	63.7–72.2
Dental insurance					
Insured	% of people	74.2	74.5	73.8	74.5
	95% CI	71.8–76.6	69.3–79.0	70.0–77.4	70.8–77.9
Uninsured	% of people	48.3	44.0	53.7	48.2
	95% CI	46.0–50.7	39.0–49.1	49.8–57.5	43.9–52.5

(a) 95% CI = 95% confidence interval for estimated percentage.

Attendance at private dental practice

While most Australians obtain dental care at private dental practices, alternatives exist in the public sector for targeted population groups. The two largest public programs are school dental services targeted to children; and adult public programs provided through dental hospitals, community health centres and regional facilities and targeted to adults holding a government concession card. In NSAOH, people were asked 'Where did you make your last dental visit?' and seven responses were offered. People who reported having visited a general dental practice, a specialist dental practice or a dental clinic associated with a health insurance fund were classified as having attended a private dental practice. In NSW, 80.8% of people aged 15 years or more attended a private practice at their last dental visit (Table 18). This estimate was not significantly different from the national estimate of 83.1% (Slade et al. 2007).

Key findings

- Adults aged 15–34 years were less likely to have visited a private dental practice at their last dental visit (74.4%) than older adults.
- Males and females were equally likely to have visited a private practice at their last dental visit irrespective of age.
- Sydney residents were more likely to attend a private practice than other residents of NSW but the differences were not statistically significant (82.0% versus 78.6%).
- The percentage was higher among people living in high socioeconomic postcodes (86.2%) than low socioeconomic postcodes (74.6%). This pattern was evident in all age groups.
- Despite having a government health card, 65.0% of cardholders reported they visited a private practice at their last dental visit, with those aged 55 years or more having the highest percentage (71.4%).
- Insured people were more likely to have visited a private practice at their last dental visit than those without dental insurance (88.0% versus 74.4%), with differences being evident in all age groups.

Discussion

The majority of NSW residents visited a private practice at their last dental visit. Insured adults and those living in areas of high socioeconomic status were more likely to do so than other population groups. Young adults were the least likely age group to report attendance at private dental practice, which may be due to people in this age group having visited the school dental service. Despite having a government health card, 65.0% of cardholders last attended a private practice, with those aged 55 years or more having the highest percentage.

In summary, socioeconomic status was only moderately associated with private visiting. This is most likely due to adults with a government health card electing to attend a private practice due to the long public waiting lists.

Table 18: Percentage of people who attended a private dental practice at last dental visit

		Population: all people Age (years)			
		All ages	15–34	35–54	≥55
All people	Per cent of people	80.8	74.4	85.9	81.9
	95% CI ^(a)	78.6–82.7	70.4–78.0	83.3–88.1	79.2–84.3
Sex					
Males	% of people	80.5	74.8	83.6	83.5
	95% CI	77.6–83.1	68.9–79.9	79.7–86.8	80.0–86.6
Females	% of people	81.0	74.1	88.2	80.4
	95% CI	78.6–83.2	68.9–78.6	85.4–90.5	77.2–83.3
Residential location					
Capital city	% of people	82.0	76.2	86.4	84.1
	95% CI	79.3–84.5	71.1–80.6	83.1–89.1	80.7–87.1
Other places	% of people	78.6	70.8	85.1	78.9
	95% CI	74.9–81.8	64.0–76.7	80.5–88.7	74.4–82.7
Postcode socioeconomic status					
Lowest	% of people	74.6	63.6	81.8	77.0
	95% CI	70.5–78.3	55.5–71.0	76.5–86.1	72.6–80.8
Middle	% of people	81.2	74.6	86.4	83.4
	95% CI	77.8–84.2	68.5–79.8	82.4–89.7	78.9–87.1
Highest	% of people	86.2	83.3	89.1	86.0
	95% CI	83.2–88.7	77.5–87.9	84.8–92.3	81.1–89.8
Government health card					
Health care card or pensioner concession card	% of people	65.0	53.8	58.3	71.4
	95% CI	61.2–68.7	44.0–63.3	51.1–65.2	67.3–75.2
Neither card	% of people	86.3	78.5	90.6	92.1
	95% CI	84.2–88.1	74.2–82.2	88.5–92.4	89.7–94.0
Dental insurance					
Insured	% of people	88.0	83.0	90.4	90.2
	95% CI	85.7–90.0	77.3–87.6	87.8–92.5	86.8–92.9
Uninsured	% of people	74.4	67.5	80.6	75.9
	95% CI	71.3–77.2	62.0–72.5	76.3–84.2	72.5–79.0

(a) 95% CI = 95% confidence interval for estimated percentage.

Payments by patients for dental care

While the place of the last visit was dominated by private practice, some visits made to private dentists are paid for by public funds. In order to identify such visits, NSAOH participants who had a government health card and who had visited a dentist within the last 5 years were asked 'Did the government or an insurance fund pay any part of the expense for your last dental visit?'. A number of response options were available including 'Paid all own expenses', 'Insurance paid some – patient paid some', 'Insurance paid all', 'Government paid some – patient paid some' and 'Government paid all'. People who reported one of the first three payment mechanisms were classified as having paid for their care, together with people who did not have a government health card and who had visited within the last 5 years. In NSW, 92.3% of people aged 15 years or more who had seen a dentist within the preceding 5 years paid for that visit (Table 19). This estimate was not significantly different from the national estimate of 91.4% (Slade et al. 2007).

Key findings

- NSW residents aged 55 years or more were less likely to have paid for their last dental visit (87.9%) than residents aged less than 55 years (94%).
- The percentages were similar for males and females (92.6% versus 92.0%).
- Sydney residents were more likely to report they had paid for their last dental visit than other residents of NSW (93.9% versus 89.5%) although the difference was fairly small. Similarly, within each age group, the percentage was higher Sydney for residents than other places, but the differences were small and not statistically significant.
- Residents of high socioeconomic postcodes were more likely to have paid for their last dental visit (96.0%) than those living in low socioeconomic postcodes (86.5%), and this pattern was consistent within all age groups.
- Despite having a government health card, 67.4% of cardholders who visited a dentist within the preceding 5 years paid for their last dental visit. Those aged 55 years or more had the highest percentage (73.7%), which may be due to older adults electing to attend a private practice due to long waiting lists in public dental care.
- Almost 100% of people with dental insurance paid for their last dental visit compared with 85.2% of uninsured people. Within the uninsured population, those aged 55 years or more reported the lowest percentage (79.5%).

Discussion

More than 9 out of 10 NSW residents who had seen a dentist in the preceding 5 years paid for that visit. Despite having a government health card, 67.4% of cardholders paid for their last dental visit.

Table 19: Percentage of people who paid for their last dental visit

		Population: people who visited dentist within last 5 years			
		Age (years)			
		All ages	15–34	35–54	≥55
All people	Per cent of people	92.3	94.4	93.8	87.9
	95% CI ^(a)	91.0–93.5	92.4–96.0	91.8–95.4	85.5–89.9
Sex					
Males	% of people	92.6	96.1	93.2	87.8
	95% CI	90.6–94.2	92.7–97.9	90.1–95.3	84.3–90.6
Females	% of people	92.0	92.9	94.5	88.0
	95% CI	90.5–93.3	89.8–95.2	92.3–96.1	85.1–90.4
Residential location					
Capital city	% of people	93.9	96.1	95.2	89.2
	95% CI	92.4–95.1	93.5–97.6	92.8–96.8	86.1–91.6
Other places	% of people	89.5	91.1	91.3	86.0
	95% CI	86.7–91.8	86.6–94.2	87.1–94.3	81.9–89.3
Postcode socioeconomic status					
Lowest	% of people	86.5	88.8	88.4	82.3
	95% CI	83.6–89.0	83.1–92.7	83.3–92.1	77.7–86.0
Middle	% of people	93.7	95.2	95.4	89.1
	95% CI	91.9–95.1	92.1–97.1	92.5–97.3	85.6–91.9
Highest	% of people	96.0	98.0	97.0	92.2
	95% CI	94.5–97.0	95.0–99.3	94.7–98.3	88.5–94.8
Government health card					
Health care card or pensioner concession card	% of people	67.4	62.4	56.0	73.7
	95% CI	63.2–71.3	51.4–72.2	47.3–64.4	69.3–77.6
Neither card	% of people	100.0	100.0	100.0	100.0
	95% CI	—	—	—	—
Place of last dental visit					
Cardholder/Public	% of people	7.3	15.7	4.1	2.6
	95% CI	3.5–14.7	5.6–36.8	1.4–11.2	0.7–9.7
Cardholder/Non-public	% of people	90.8	96.5	87.7	90.2
	95% CI	88.1–92.9	88.9–99.0	78.1–93.4	87.1–92.6
Non-cardholder/Non-public	95% CI	100.0	100.0	100.0	100.0
	% of people	—	—	—	—
Dental insurance					
Insured	% of people	99.1	99.4	99.5	98.1
	95% CI	98.3–99.5	95.6–99.9	98.3–99.8	96.3–99.0
Uninsured	% of people	85.2	89.6	86.3	79.5
	95% CI	82.8–87.4	85.7–92.5	82.1–89.7	75.9–82.7

(a) 95% CI = 95% confidence interval for estimated percentage.

Government-subsidised dental care in private sector

In some states and territories, public sector dental programs provide care to people eligible for their services by referring them to private practitioner dentists. The cost of such care is then subsidised by the state or territory dental program. In NSW, 1.6% of the adult population received state-subsidised dental care in the private sector (Table 20). This statistic was not reported nationally.

- People aged 55 years or more were significantly more likely than people in the two younger age groups to receive state-subsidised dental care in private practice.
- However, the age-related pattern was not statistically significant within the group of people who had a government health card, among whom 6.9% received state-subsidised dental care in private practice.
- Dental insurance status was the only characteristic associated with a significant difference in likelihood of state-subsidised dental care in private practice.

Discussion

Variation in this statistic according to age and dental insurance status reflected similar variation in the distribution of people who were eligible for state dental services. When the data were limited to people eligible for state dental services, there were no meaningful differences among age groups.

Table 20: Percentage of people who received government-subsidised dental care in private sector

		Population: people who visited dentist within last 5 years			
		Age (years)			
		All ages	15–34	35–54	≥55
All people	Per cent of people	1.6	0.8	1.0	3.4
	<i>95% CI^(a)</i>	<i>1.3–2.1</i>	<i>0.4–1.6</i>	<i>0.5–2.0</i>	<i>2.5–4.6</i>
Sex					
Males	% of people	1.3	0.5	1.1	2.5
	<i>95% CI</i>	<i>0.8–2.0</i>	<i>0.1–2.1</i>	<i>0.4–2.8</i>	<i>1.4–4.4</i>
Females	% of people	2.0	1.0	1.0	4.2
	<i>95% CI</i>	<i>1.4–2.7</i>	<i>0.4–2.5</i>	<i>0.4–2.5</i>	<i>2.9–6.1</i>
Residential location					
Capital city	% of people	1.4	0.6	0.8	3.4
	<i>95% CI</i>	<i>1.0–2.0</i>	<i>0.2–1.6</i>	<i>0.3–1.9</i>	<i>2.2–5.1</i>
Other places	% of people	2.0	1.1	1.5	3.5
	<i>95% CI</i>	<i>1.4–3.0</i>	<i>0.4–3.7</i>	<i>0.6–3.8</i>	<i>2.2–5.5</i>
Postcode socioeconomic status					
Lowest	% of people	2.5	1.4	2.3	4.0
	<i>95% CI</i>	<i>1.8–3.6</i>	<i>0.4–4.3</i>	<i>1.1–4.7</i>	<i>2.5–6.1</i>
Middle	% of people	1.0	0.2	0.4	3.2
	<i>95% CI</i>	<i>0.6–1.7</i>	<i>0.0–1.3</i>	<i>0.1–1.2</i>	<i>1.8–5.5</i>
Highest	% of people	1.4	0.9	0.6	3.1
	<i>95% CI</i>	<i>0.9–2.3</i>	<i>0.3–2.8</i>	<i>0.1–3.8</i>	<i>1.7–5.7</i>
Government health card					
Health care card or pensioner concession card	% of people	6.9	5.1	7.2	7.4
	<i>95% CI</i>	<i>5.3–8.9</i>	<i>2.4–10.8</i>	<i>3.8–13.5</i>	<i>5.6–9.9</i>
Neither card	% of people	0.0	0.0	0.0	0.0
	<i>95% CI</i>	—	—	—	—
Dental insurance					
Insured	% of people	0.3	0.0	0.2	0.7
	<i>95% CI</i>	<i>0.1–0.8</i>	—	<i>0.0–1.2</i>	<i>0.3–1.9</i>
Uninsured	% of people	3.1	1.5	2.2	5.7
	<i>95% CI</i>	<i>2.3–4.0</i>	<i>0.7–3.2</i>	<i>1.0–4.4</i>	<i>4.1–7.7</i>

(a) 95% CI = 95% confidence interval for estimated percentage.

People's usual pattern of dental visits

While time since last visiting a dentist provides a snapshot of dental visiting behaviour, people's usual dental attendance pattern reflects longer term behaviours and intentions. In NSAOH, people who were dentate were asked 'How often on average do you seek care from a dental professional?' and four categories of response were offered. In NSW, 54.9% of people aged 15 years or more usually visit a dentist at least once a year (Table 21). This estimate was not significantly different from the national estimate of 53.1% (Slade et al. 2007).

Key findings

- Dental visiting behaviour was very similar across age groups, with 53.8% of young adults usually visiting at least once a year compared with 56.7% of adults aged 55 years or more.
- Females were more likely than males to usually visit a dentist at least once a year (60.8% versus 49.0%). This pattern was consistent across all age groups although differences were not significant for the 15–34 years age group.
- The percentage was higher for Sydney residents than people living in other places (58.8% versus 47.8%). This difference was mainly attributable to those aged 55 years or more, with 63.2% of Sydney residents usually visiting at least once a year compared with 47.1% of other residents of NSW.
- Residents of high socioeconomic postcodes were much more likely to usually visit one or more times a year than those in low socioeconomic postcodes (64.0% versus 48.0%). Differences were mainly evident in the two older age groups.
- People who were government health cardholders were less likely to attend annually than non-government health cardholders (44.4% versus 58.1%). This pattern was consistent across all age groups although differences were not significant for the 15–34 years age group.
- Among government health cardholders, those who had visited a private practice at their last dental visit were more likely to attend annually than those who last visited a public practice (47.8% versus 35.6%). Although this difference was not statistically significant, those holding a government health card aged 55 years or more who had attended a private practice were twice as likely to report usually visiting at least once a year than those who attended a public practice.
- Insured people were far more likely to usually visit a dentist one or more times a year than uninsured people (69.3% versus 41.1%), with large differences being evident in all age groups.

Discussion

Just over half of NSW residents aged 15 years or more usually visit the dentist at least once a year. Being female, residing in the metropolitan region, living in areas of high socioeconomic status, being a non-government health cardholder and having dental insurance were all associated with regular dental visiting. Differences in visiting behaviour between population groups were most evident between insured and uninsured groups. Comparisons by age highlighted larger differences between population groups for adults aged 35–54 years and 55 years or more.

Table 21: Percentage of people who usually visit a dental professional at least once a year

		Population: dentate people Age (years)			
		All ages	15–34	35–54	≥55
All people	Per cent of people	54.9	53.8	54.5	56.7
	95% CI ^(a)	52.5–57.2	49.6–58.0	51.0–58.0	53.1–60.2
Sex					
Males	% of people	49.0	49.4	47.5	50.3
	95% CI	45.7–52.2	42.6–56.3	43.0–52.0	45.5–55.2
Females	% of people	60.8	58.3	61.6	62.8
	95% CI	57.8–63.7	53.3–63.1	57.3–65.8	58.8–66.7
Residential location					
Capital city	% of people	58.8	56.3	58.5	63.2
	95% CI	55.6–62.0	51.2–61.2	53.8–63.1	58.4–67.8
Other places	% of people	47.8	48.8	47.5	47.1
	95% CI	44.7–50.9	41.5–56.2	42.7–52.3	41.7–52.6
Postcode socioeconomic status					
Lowest	% of people	48.0	52.5	46.6	44.8
	95% CI	44.5–51.6	46.7–58.4	40.5–52.8	38.7–51.1
Middle	% of people	51.9	50.6	50.3	56.3
	95% CI	47.7–56.0	43.0–58.2	45.1–55.5	49.9–62.4
Highest	% of people	64.0	58.4	66.1	68.6
	95% CI	60.4–67.5	50.7–65.7	61.0–70.9	63.6–73.1
Government health card					
Health care card or pensioner concession card	% of people	44.4	43.6	39.5	46.8
	95% CI	40.3–48.5	34.7–53.1	32.7–46.9	41.9–51.8
Neither card	% of people	58.1	56.1	57.0	64.5
	95% CI	55.4–60.7	51.5–60.6	53.2–60.7	59.9–68.8
Place of last dental visit					
Cardholder/Public	% of people	35.6	47.5	32.7	23.8
	95% CI	28.1–43.9	32.5–63.0	21.8–45.9	15.9–34.1
Cardholder/Non-public	% of people	47.8	40.6	43.5	51.4
	95% CI	43.0–52.6	28.5–54.0	34.7–52.9	45.6–57.2
Non-cardholder/Non-public	% of people	58.1	56.1	57.0	64.5
	95% CI	55.4–60.7	51.5–60.6	53.2–60.7	59.9–68.8
Dental insurance					
Insured	% of people	69.3	70.3	67.4	71.2
	95% CI	66.2–72.3	63.3–76.5	63.0–71.5	66.9–75.1
Uninsured	% of people	41.1	40.3	38.8	44.6
	95% CI	38.4–43.8	35.4–45.3	34.5–43.2	40.6–48.8

(a) 95% CI = 95% confidence interval for estimated percentage.

Usual attendance at the same dentist

In NSAOH, usual source of care was assessed in the interview by asking people 'Is there a dentist you usually go to for dental care?'. People who answered 'yes', have a usual source of care' were classified as having a dentist they usually attend. In NSW, 81.4% of the dentate population aged 15 years or more who visited a dentist within the last 5 years reported having a dentist they usually attend (Table 22), which was slightly higher, but not significantly, than the national estimate of 78.6% (Slade et al. 2007).

Key findings

- Across age groups the percentage who usually attended the same dentist increased from 74.3% among 15–34-year-olds to 87.0% among people aged 55 years or more.
- A greater percentage of females than males reported having a dentist they usually attend, both for all ages combined and across all age groups.
- For people of all ages there was less variation among groups classified by residential location. Percentages were higher for residents living in the capital city compared with other places (82.9% versus 78.8%).
- People living in postcodes with low socioeconomic status were less likely to report a usual source of care compared with those in postcodes with high socioeconomic status (76.2% versus 86.2%). Significant age-specific differences were found by low and high socioeconomic status in the 15–34 years (67.2% versus 81.1%) and 35–54 years (79.2% versus 88.3%) age groups.
- Among non-government health cardholders, a significantly higher percentage reported having a dentist they usually attend than among government health cardholders (83.4% versus 74.8%). The largest difference occurred in the 35–54 years age group (86.6% versus 68.0%).
- Within the population of government health cardholders, people whose last dental visit was to the public sector were less likely to report having a dentist they usually attend than those who attended elsewhere (54.7% versus 82.6%). The largest differences occurred in the 35–54 years (45.2% versus 81.3%) and 55 years or more (51.0% versus 84.6%) age groups. The percentage was lower for people with a government health card who last visited the public sector than those who were non-government health cardholders and attended elsewhere (54.7% versus 83.4%).
- The percentage was significantly higher among adults with dental insurance than for those uninsured (90.2% versus 72.3%). This pattern was consistent across all age groups.

Discussion

In summary, 81.4% of NSW adults reported that they usually visit the same dentist. This type of visiting was more frequent among the older age groups, females and those who are insured.

Table 22: Percentage of people who have a dentist they usually attend

		Population: dentate people who visited dentist within last 5 years			
		Age (years)			
		All ages	15–34	35–54	≥55
All people	Per cent of people	81.4	74.3	84.0	87.0
	95% CI ^(a)	79.3–83.3	70.4–77.9	81.2–86.5	84.5–89.3
Sex					
Males	% of people	78.7	71.3	80.1	86.5
	95% CI	75.7–81.4	64.9–77.0	75.5–84.1	82.3–89.8
Females	% of people	83.9	77.1	87.8	87.6
	95% CI	81.6–86.0	72.7–81.0	84.9–90.2	84.4–90.2
Residential location					
Capital city	% of people	82.9	76.9	84.7	88.8
	95% CI	80.5–85.0	72.3–81.0	81.2–87.7	85.6–91.3
Other places	% of people	78.8	69.0	82.8	84.4
	95% CI	74.8–82.2	61.4–75.6	77.7–86.9	79.5–88.2
Postcode socioeconomic status					
Lowest	% of people	76.2	67.2	79.2	82.3
	95% CI	72.4–79.7	59.1–74.3	73.9–83.6	78.4–85.7
Middle	% of people	80.9	73.2	84.0	88.9
	95% CI	77.3–84.1	66.7–78.9	79.1–87.9	83.5–92.7
Highest	% of people	86.2	81.1	88.3	89.6
	95% CI	83.3–88.7	75.1–86.0	83.8–91.7	85.0–92.9
Government health card					
Health care card or pensioner concession card	% of people	74.8	72.9	68.0	78.6
	95% CI	71.1–78.2	64.5–79.9	59.3–75.6	74.3–82.4
Neither card	% of people	83.4	74.9	86.6	93.4
	95% CI	81.1–85.5	70.4–78.9	83.7–89.0	90.3–95.5
Place of last dental visit					
Cardholder/Public	% of people	54.7	66.1	45.2	51.0
	95% CI	47.1–62.1	49.4–79.5	30.7–60.5	40.5–61.4
Cardholder/Non-public	% of people	82.6	78.0	81.3	84.6
	95% CI	78.8–85.9	67.8–85.6	72.3–87.9	79.7–88.4
Non-cardholder/Non-public	% of people	83.4	74.9	86.6	93.4
	95% CI	81.1–85.5	70.4–78.9	83.7–89.0	90.3–95.5
Dental insurance					
Insured	% of people	90.2	85.9	90.6	95.1
	95% CI	88.3–91.8	81.3–89.5	87.6–92.9	92.5–96.8
Uninsured	% of people	72.3	64.0	75.0	79.7
	95% CI	69.1–75.2	58.3–69.3	70.3–79.2	75.9–83.0

(a) 95% CI = 95% confidence interval for estimated percentage.

Usual dental attendance for a check-up

In NSAOH, dentate people were asked 'Is your usual reason for visiting a dental professional for check-ups or when you have a dental problem?'. In NSW, 56.8% of the adult dentate population reported usually visiting a dentist for a check-up (Table 23), which was not significantly different from the national estimate of 56.2% (Slade et al. 2007).

Key findings

- Across age groups there was little variation in the percentage of adults usually visiting a dentist for a check-up.
- The percentage of adults was significantly higher among females than males (62.1% versus 51.5%). This difference was mainly attributable to those aged 35–54 years, with 61.8% of females reporting usually visiting for a check-up compared with only 48.5% of males.
- The percentage was higher for people living in Sydney than the rest of NSW (61.1% versus 49.2%). This difference was mainly attributable to those aged 55 years or more, with 62.5% of Sydney residents reporting usually visiting for a check-up compared with only 45.3% of those living outside of Sydney.
- For each age group, the percentage was significantly lower for adults living in low compared with high socioeconomic postcodes. In those aged 55 years or more, 46.6% with low socioeconomic status reported usually visiting for a check-up compared with those with middle (54.4%) and high (68.5%) socioeconomic status.
- The percentage was significantly lower for adults who had a government health card than for those who did not (40.4% versus 61.6%). This pattern was consistent across all age groups.
- Within the population of government health cardholders, people whose last dental visit was to the public sector were less likely to report usually visiting a dentist for a check-up than those who attended elsewhere (23.3% versus 47.0%). The largest differences occurred in the 35–54 years (9.1% versus 39.8%) and the 55 years or more (14.2% versus 48.3%) age groups.
- The percentage was significantly higher among adults with dental insurance than for those uninsured (71.3% versus 42.6%). This pattern was consistent across all age groups.

Discussion

In summary, just over half of the adult population usually visited the dentist for a check-up, with this percentage being slightly higher for those aged 15–34 years. There was significant association with being female, living in Sydney, living in a high socioeconomic postcode, not having a government health card and having dental insurance. The percentage was markedly more frequent among non-government health cardholders and those with dental insurance.

Table 23: Percentage of people who usually visit a dentist for a check-up

		Population: dentate people Age (years)			
		All ages	15–34	35–54	≥55
All people	Per cent of people	56.8	59.5	55.2	55.5
	95% CI ^(a)	54.4–59.2	55.3–63.6	51.5–58.8	52.0–58.9
Sex					
Males	% of people	51.5	54.3	48.5	51.8
	95% CI	48.2–54.7	47.5–60.8	43.8–53.2	47.3–56.3
Females	% of people	62.1	64.8	61.8	59.1
	95% CI	59.1–65.0	59.8–69.4	57.3–66.1	54.9–63.1
Residential location					
Capital city	% of people	61.1	62.8	58.5	62.5
	95% CI	57.9–64.3	57.4–68.0	53.9–63.0	57.7–67.0
Other places	% of people	49.2	52.8	49.2	45.3
	95% CI	45.6–52.7	45.9–59.5	43.4–55.0	40.5–50.1
Postcode socioeconomic status					
Lowest	% of people	46.6	51.5	45.9	42.0
	95% CI	43.4–49.9	44.3–58.6	40.5–51.5	38.1–46.1
Middle	% of people	54.4	57.7	50.9	54.4
	95% CI	50.2–58.5	50.3–64.8	44.8–57.0	48.2–60.5
Highest	% of people	68.5	68.2	67.8	69.6
	95% CI	65.2–71.5	61.0–74.7	62.7–72.6	64.9–73.9
Government health card					
Health care card or pensioner concession card	% of people	40.4	46.8	28.4	42.5
	95% CI	36.7–44.1	37.6–56.2	21.2–37.0	38.3–46.9
Neither card	% of people	61.6	61.7	59.5	65.8
	95% CI	59.1–64.1	57.3–66.0	55.8–63.1	61.1–70.1
Place of last dental visit					
Cardholder/Public	% of people	23.3	42.4	9.1	14.2
	95% CI	16.8–31.3	27.9–58.3	3.9–19.5	8.6–22.5
Cardholder/Non-public	% of people	47.0	50.2	39.8	48.3
	95% CI	43.0–51.0	37.6–62.8	30.1–50.4	43.3–53.3
Non-cardholder/Non-public	% of people	61.6	61.7	59.5	65.8
	95% CI	59.1–64.1	57.3–66.0	55.8–63.1	61.1–70.1
Dental insurance					
Insured	% of people	71.3	75.4	68.1	71.5
	95% CI	68.3–74.1	68.8–80.9	63.8–72.0	66.8–75.8
Uninsured	% of people	42.6	45.6	39.4	42.3
	95% CI	39.7–45.6	40.0–51.4	34.6–44.3	38.5–46.1

(a) 95% CI = 95% confidence interval for estimated percentage.

Dental care avoided or delayed due to cost

In NSAOH, cost as a barrier to receipt of dental care was assessed with the question 'During the last 12 months, have you avoided or delayed visiting a dental professional because of the cost?'. People who answered 'yes' were classified as having delayed or avoided dental care due to cost. In NSW, they represented 29.8% of the population aged 15 years or more (Table 24), which was not significantly different from the national estimate of 30.0% (Slade et al. 2007).

Key findings

- The percentage reporting cost as a barrier to receipt of dental care was lower among people aged 55 years or more (23.7%) compared with younger age groups.
- Females were more likely than males to have avoided or delayed care due to cost (32.7% versus 26.9%). This pattern was consistent across all age groups although the differences were not statistically significant.
- In the two older age groups, the percentage was lower among people living in Sydney than those living outside the capital city. The reverse was true for the younger age group although this difference was not statistically significant.
- People living in postcodes with low socioeconomic status were more likely to report avoidance or delay than those in postcodes with high socioeconomic status (34.7% versus 26.4%). Statistically significant differences were observed in the 35–54 years (40.6% versus 26.0%) and 55 years or more (31.3% versus 15.9%) age groups.
- Avoiding or delaying care due to cost occurred more frequently among adults who had a government health card than for those who did not (39.1% versus 26.8%). This pattern was consistent across all age groups, with the largest difference observed in the 35–54 years age group.
- Within the population of government health cardholders, the percentage tended to be higher among people whose last dental visit was to the public sector than for those who attended elsewhere (44.2% versus 37.1%). However, the pattern was not consistent within age groups and 95% CIs were large; hence, the differences were not statistically significant.
- Uninsured adults were more likely to report this cost barrier than insured adults, a pattern that was consistent across all age groups. For each age group, there was over a two-fold difference between the insured and uninsured.

Discussion

In summary, dental insurance was strongly associated with having avoided or delayed receipt of dental care due to cost. There was a moderate association with sex, postcode socioeconomic status and government health cardholder status.

Table 24: Percentage of people who avoided or delayed dental care

		Population: all people Age (years)			
		All ages	15–34	35–54	≥55
All people	Per cent of people	29.8	32.0	32.9	23.7
	95% CI ^(a)	27.9–31.8	28.6–35.7	29.9–36.1	21.4–26.3
Sex					
Males	% of people	26.9	27.1	30.7	21.9
	95% CI	24.1–29.8	22.0–33.0	26.2–35.6	18.6–25.6
Females	% of people	32.7	37.0	35.1	25.4
	95% CI	30.1–35.4	32.1–42.2	31.4–39.0	22.6–28.5
Residential location					
Capital city	% of people	27.8	32.6	29.4	19.4
	95% CI	25.4–30.3	28.2–37.2	25.9–33.2	16.4–22.7
Other places	% of people	33.2	30.9	38.9	29.6
	95% CI	30.0–36.7	25.6–36.8	33.5–44.7	26.1–33.4
Postcode socioeconomic status					
Lowest	% of people	34.7	32.0	40.6	31.3
	95% CI	31.4–38.2	25.5–39.3	34.9–46.6	27.8–35.1
Middle	% of people	28.6	28.9	32.6	23.0
	95% CI	25.5–31.9	23.8–34.5	27.9–37.6	18.8–27.8
Highest	% of people	26.4	35.5	26.0	15.9
	95% CI	23.2–29.9	29.6–41.8	21.5–31.2	12.6–19.8
Government health card					
Health care card or pensioner concession card	% of people	39.1	46.0	54.9	31.1
	95% CI	35.4–42.9	37.2–55.0	47.3–62.3	27.7–34.8
Neither card	% of people	26.8	29.8	29.2	16.7
	95% CI	24.8–29.0	26.2–33.7	26.1–32.4	14.2–19.6
Place of last dental visit					
Cardholder/Public	% of people	44.2	37.5	56.7	40.9
	95% CI	37.4–51.2	24.1–53.1	43.6–69.0	32.9–49.3
Cardholder/Non-public	% of people	37.1	52.9	53.8	29.0
	95% CI	33.1–41.3	40.3–65.1	43.9–63.4	25.3–33.0
Non-cardholder/Non-public	% of people	26.8	29.8	29.2	16.7
	95% CI	24.8–29.0	26.2–33.7	26.1–32.4	14.2–19.6
Dental insurance					
Insured	% of people	17.4	17.5	20.0	13.5
	95% CI	15.5–19.6	13.5–22.5	17.0–23.3	10.7–16.9
Uninsured	% of people	41.4	45.1	48.7	31.0
	95% CI	38.7–44.2	40.2–50.1	43.8–53.6	27.8–34.4

(a) 95% CI = 95% confidence interval for estimated percentage.

Recommended dental treatment foregone due to cost

In NSAOH, treatment foregone due to cost was assessed with the question 'Has the cost prevented you from having any dental treatment that was recommended during the last 2 years?'. People who answered 'yes' were classified as having foregone dental treatment due to cost. In NSW, they represented 20.0% of the population aged 15 years or more (Table 25), which was slightly lower, but not significantly, than the national estimate of 20.6% (Slade et al. 2007).

Key findings

- Across age groups, there was some variation in frequency of foregone treatment, with 24.6% of adults aged 35–54 years reporting that they had forgone treatment due to cost compared with 14.8% of those aged 55 years or more.
- For all ages combined, the percentage was higher in females compared with males (21.4% versus 18.5%). This difference, however, was not statistically significant. For each age group, there was no statistically significant difference in the percentage of males and females not having recommended dental treatment.
- For people of all ages, there was less variation among groups classified by residential location. Percentages were lower for residents in the capital city compared with other places (19.3% versus 21.5%). However, no statistically significant differences were found between people living in Sydney and those living outside the capital city within each age group.
- People living in postcodes with low socioeconomic status were significantly more likely to report forgoing recommended dental treatment due to cost than those in postcodes with high socioeconomic status (24.3% versus 16.6%). This pattern was consistent across all age groups although the differences were not statistically significant.
- The percentage was significantly higher for adults who were government health cardholders than for those who were not (29.2% versus 17.4%). The largest differences occurred in the 15–34 years (33.8% versus 16.9%) and 35–54 years (50.7% versus 20.7%) age groups.
- There was some variation according to place of last visit. For example, within the population of government health cardholders, people whose last dental visit was to the public sector were more likely to report forgoing recommended dental treatment due to cost than those who attended elsewhere (41.4% versus 24.7%). This pattern was consistent across all age groups but, because 95% CIs were large in these groups, the differences were not statistically significant.
- The percentage was significantly higher among adults with no dental insurance than for those uninsured (28.5% versus 13.0%). This pattern was consistent across all age groups.

Discussion

In summary, having foregone recommended dental treatment due to cost was strongly associated with dental insurance. There was a moderate association with postcode socioeconomic status and government health cardholder status.

Table 25: Percentage of people who reported that cost had prevented recommended dental treatment

		Population: people who visited dentist within last 2 years			
		Age (years)			
		All ages	15–34	35–54	≥55
All people	Per cent of people	20.0	19.4	24.6	14.8
	95% CI ^(a)	18.0–22.2	15.7–23.7	21.6–27.9	12.3–17.6
Sex					
Males	% of people	18.5	16.1	24.9	12.4
	95% CI	15.7–21.7	10.9–23.1	20.0–30.6	9.6–15.8
Females	% of people	21.4	22.1	24.3	16.8
	95% CI	19.0–24.0	17.3–27.8	21.4–27.5	13.7–20.4
Residential location					
Capital city	% of people	19.3	18.2	24.5	13.1
	95% CI	16.8–22.0	14.1–23.2	21.0–28.4	10.4–16.5
Other places	% of people	21.5	21.8	24.8	17.4
	95% CI	18.2–25.3	14.9–30.6	19.6–30.8	13.0–22.7
Postcode socioeconomic status					
Lowest	% of people	24.3	25.4	28.7	17.7
	95% CI	20.7–28.4	18.6–33.7	22.1–36.3	13.7–22.6
Middle	% of people	20.0	18.6	23.4	17.2
	95% CI	17.0–23.5	12.8–26.2	19.4–27.9	12.4–23.3
Highest	% of people	16.6	15.5	22.3	10.4
	95% CI	13.5–20.3	10.3–22.7	18.1–27.2	7.3–14.5
Government health card					
Health care card or pensioner concession card	% of people	29.2	33.8	50.7	18.7
	95% CI	24.8–34.1	22.8–46.9	42.2–59.2	14.6–23.7
Neither card	% of people	17.4	16.9	20.7	11.8
	95% CI	15.4–19.6	13.1–21.4	18.0–23.6	9.3–14.8
Place of last dental visit					
Cardholder/Public	% of people	41.4	36.9	64.5	25.6
	95% CI	33.8–49.5	20.5–57.0	50.3–76.5	16.9–36.6
Cardholder/Non-public	% of people	24.7	31.5	42.7	17.3
	95% CI	20.1–30.0	19.5–46.6	32.7–53.4	13.1–22.7
Non-cardholder/Non-public	% of people	17.4	16.9	20.7	11.8
	95% CI	15.4–19.6	13.1–21.4	18.0–23.6	9.3–14.8
Dental insurance					
Insured	% of people	13.0	11.7	15.7	10.3
	95% CI	10.9–15.5	7.9–16.9	12.7–19.3	7.4–14.2
Uninsured	% of people	28.5	28.3	38.0	18.9
	95% CI	25.3–32.0	22.3–35.3	32.2–44.2	15.2–23.2

(a) 95% CI = 95% confidence interval for estimated percentage.

Difficulty paying a \$100 dental bill

In NSAOH, difficulty paying for dental care was assessed with the question 'At most times of the year, how much difficulty would you have paying a \$100 dental bill? Would you say none, hardly any, a little, a lot of difficulty, don't know?'. People who answered 'a lot' were classified as having difficulty paying a \$100 dental bill. They represented 19.3% of the NSW population aged 15 years or more (Table 26), which was slightly higher, but not significantly, than the national estimate of 18.2% (Slade et al. 2007).

Key findings

- Across age groups there was little variation in the percentage of adults who reported difficulty paying a \$100 dental bill.
- For all ages combined, there was a significantly greater percentage of females compared with males (22.0% versus 16.5%). This difference was mainly attributable to those aged 15–34 years (27.0% of females compared with only 16.2% of males).
- The percentage was higher among people living outside Sydney than those in Sydney (22.5% versus 17.4%). This pattern was consistent across all age groups although the differences were not statistically significant.
- People living in postcodes with low socioeconomic status were significantly more likely to report that they would have difficulty paying a \$100 dental bill than those in postcodes with high socioeconomic status (23.6% versus 14.7%). Statistically significant differences were observed in the 35–54 years (23.1% versus 14.2%) and 55 years or more (22.8% versus 12.5%) age groups.
- The percentage was higher among adults who held a government health card compared with those who did not (36.1% versus 13.3%). This pattern was consistent across all age groups, and the largest difference was observed in the 35–54 years age group.
- Within the population of government health cardholders, people whose last dental visit was to the public sector were more likely to have difficulty paying a \$100 dental bill than those who attended elsewhere (52.6% versus 29.8%). Statistically significant differences occurred in the 35–54 years (68.6% versus 42.2%) and 55 years or more (50.2% versus 24.4%) age groups.
- The percentage was much higher among people with no dental insurance than the insured (26.2% versus 10.9%). This pattern was consistent across all age groups. The largest difference occurred among those aged 55 years or more, with 26.7% of uninsured adults reporting difficulty compared with 6.7% of insured adults.

Discussion

In summary, government health cardholder status and dental insurance were strongly associated with having a lot of difficulty paying a \$100 dental bill. There was a moderate association with sex, postcode socioeconomic status and place of last dental visit.

Table 26: Percentage of people who would have a lot of difficulty paying a \$100 dental bill

		Population: all people			
		Age (years)			
		All ages	15–34	35–54	≥55
All people	Per cent of people	19.3	21.5	17.8	18.4
	95% CI ^(a)	17.7–21.0	18.4–25.0	15.5–20.4	16.0–21.1
Sex					
Males	% of people	16.5	16.2	17.3	15.9
	95% CI	14.3–18.9	12.3–21.1	13.6–21.8	12.6–19.9
Females	% of people	22.0	27.0	18.3	20.7
	95% CI	19.8–24.3	22.7–31.7	15.8–21.1	17.6–24.1
Residential location					
Capital city	% of people	17.4	19.7	16.3	15.8
	95% CI	15.3–19.7	16.2–23.7	13.4–19.6	12.7–19.5
Other places	% of people	22.5	25.3	20.5	21.9
	95% CI	20.0–25.2	19.3–32.4	16.8–24.8	18.3–26.1
Postcode socioeconomic status					
Lowest	% of people	23.6	25.0	23.1	22.8
	95% CI	21.1–26.3	19.1–31.9	19.3–27.4	19.1–27.1
Middle	% of people	19.7	23.0	16.6	19.3
	95% CI	16.8–23.0	17.5–29.5	13.0–20.9	15.1–24.4
Highest	% of people	14.7	17.1	14.2	12.5
	95% CI	12.3–17.6	13.3–21.8	10.5–18.9	8.9–17.4
Government health card					
Health care card or pensioner concession card	% of people	36.1	39.9	52.3	29.1
	95% CI	32.4–39.9	30.1–50.5	44.3–60.2	25.5–33.1
Neither card	% of people	13.3	17.6	11.9	8.0
	95% CI	11.7–15.0	14.6–21.0	10.0–14.0	6.0–10.6
Place of last dental visit					
Cardholder/Public	% of people	52.6	41.7	68.6	50.2
	95% CI	45.9–59.1	28.1–56.6	56.7–78.5	41.4–59.1
Cardholder/Non-public	% of people	29.8	38.5	42.2	24.4
	95% CI	26.0–33.9	26.8–51.7	32.8–52.2	20.8–28.4
Non-cardholder/Non-public	% of people	13.3	17.6	11.9	8.0
	95% CI	11.7–15.0	14.6–21.0	10.0–14.0	6.0–10.6
Dental insurance					
Insured	% of people	10.9	15.4	10.2	6.7
	95% CI	9.1–12.9	11.5–20.3	8.0–12.9	4.7–9.3
Uninsured	% of people	26.2	25.1	27.1	26.7
	95% CI	24.0–28.6	20.8–29.9	22.9–31.7	23.5–30.2

(a) 95% CI = 95% confidence interval for estimated percentage.

Percentage of people avoiding foods due to dental problems

Avoiding food due to dental problems is a sign of poor oral health and may reflect an inability to eat properly. This reduces enjoyment of food and could affect the ability to maintain a healthy nutritional status.

In NSAOH, avoiding food was assessed in the interview by asking people 'How often have you had to avoid eating some foods because of problems with your teeth, mouth or dentures during the last 12 months? Was it: very often, often, sometimes, hardly ever, never during the last 12 months, don't know?'. People who answered 'very often', 'often' or 'sometimes' were classified as having avoided certain foods. They represented 19.7% of the NSW population aged 15 years or more (Table 27), which was higher than the national estimate of 17.4% (Slade et al. 2007). The difference was not statistically significant.

Key findings

- Females were more likely to report that they avoided some food (22.5%) than males (16.9%). The gap between females and males increased with age.
- There was a similar difference between those who live outside capital cities (23.6%) compared with those who live in a capital city (17.5%).
- The percentage who avoided food decreased as the socioeconomic status of the postcode increased. It was highest in the lowest socioeconomic postcode (22.7%) and lowest in the highest socioeconomic postcode (15.3%).
- Frequency of avoiding food was twice as high in people who were government health cardholders (31.9%) compared with non-government health cardholders (15.7%). The difference was greatest in the youngest age group and attenuated with age.
- Within the population of government health cardholders, those who last visited a public dental clinic were almost twice as likely (44.8%) as those who visited a private dentist (26.9%) to avoid foods.
- People with no dental insurance were more likely (24.5%) than those with insurance (14.3) to avoid foods.

Discussion

Residents of NSW were equally as likely as the rest of the Australian population to avoid some foods because of problems with their teeth, mouth or gums. Avoiding some foods because of dental problems was associated with being female, living outside a capital city or in a low socioeconomic postcode, having last visited a public clinic and not having dental insurance.

Table 27: Percentage of people avoiding foods due to dental problems

		Population: all people Age (years)			
		All ages	15–34	35–54	≥55
All people	Per cent of people	19.7	17.5	18.8	23.2
	95% CI ^(a)	18.0–21.6	14.5–21.0	16.3–21.7	20.9–25.7
Sex					
Males	% of people	16.9	15.5	16.6	18.9
	95% CI	14.6–19.4	11.4–20.8	13.0–20.9	15.7–22.6
Females	% of people	22.5	19.6	21.1	27.1
	95% CI	20.4–24.7	15.6–24.3	18.1–24.4	24.0–30.4
Residential location					
Capital city	% of people	17.5	15.1	17.0	21.1
	95% CI	15.3–19.9	11.5–19.7	14.3–20.2	18.5–24.0
Other places	% of people	23.6	22.5	22.0	26.1
	95% CI	20.7–26.7	17.7–28.2	17.2–27.7	22.3–30.3
Postcode socioeconomic status					
Lowest	% of people	22.7	19.8	22.9	25.0
	95% CI	19.5–26.2	14.7–26.0	17.5–29.4	21.1–29.4
Middle	% of people	21.3	19.2	20.3	25.4
	95% CI	18.2–24.7	14.2–25.4	16.4–24.9	21.2–30.2
Highest	% of people	15.3	13.8	13.5	19.1
	95% CI	12.8–18.1	9.2–20.3	10.7–17.0	16.1–22.5
Government health card					
Health care card or pensioner concession card	% of people	31.9	34.1	37.6	29.1
	95% CI	28.2–35.8	24.9–44.7	29.8–46.0	26.0–32.4
Neither card	% of people	15.7	14.7	15.6	17.7
	95% CI	13.9–17.6	11.8–18.0	13.1–18.4	14.5–21.3
Place of last dental visit					
Cardholder/Public	% of people	44.8	41.1	50.8	43.7
	95% CI	36.9–53.0	26.8–57.1	39.1–62.3	35.3–52.6
Cardholder/Non-public	% of people	26.9	28.5	29.2	25.9
	95% CI	23.3–30.7	17.5–42.7	20.9–39.2	22.7–29.3
Non-cardholder/Non-public	% of people	15.7	14.7	15.6	17.7
	95% CI	13.9–17.6	11.8–18.0	13.1–18.4	14.5–21.3
Dental insurance					
Insured	% of people	14.3	13.3	13.5	16.9
	95% CI	12.6–16.2	10.1–17.3	11.2–16.3	13.7–20.7
Uninsured	% of people	24.5	21.0	25.0	27.8
	95% CI	21.9–27.4	16.5–26.4	20.8–29.6	24.9–30.9

(a) 95% CI = 95% confidence interval for estimated percentage.

5 Oral health perceptions

Percentage of people rating their oral health as fair or poor

Self-reported global measures of oral health reflect an individual's own experience of their oral health. Single-item, self-rated oral health measures are associated with functional impairment and discomfort as well as clinical measures of dental health. They are used widely in research and provide a summary measure of oral symptoms and functioning (Benyamini et al. 2004).

In NSAOH, self-rated oral health was assessed in the interview by asking people 'And how would you rate your own DENTAL health. Would you say that it is: excellent, very good, good, fair, poor, don't know?'. People who answered 'fair' or 'poor' were classified as having fair or poor self-rated oral health. They represented 16.9% of the NSW population aged 15 years or more (Table 28), which is very close to the national estimate of 16.4% (Slade et al. 2007). The difference was not statistically significant.

Key findings

- The percentage who reported fair or poor oral health decreased as the socioeconomic status of the postcode where they lived increased. It was highest in the lowest socioeconomic postcode (21.1%) and lowest in the highest socioeconomic postcode (14.4%).
- The percentage reporting fair or poor health was more than twice as high in government health cardholders (29.6%) than non-government health cardholders (13.2%). The difference was larger in the 15–34 years and 35–44 years age groups than in the 55 years or more age group.
- Within the population of government health cardholders, those who last visited a public dental clinic were almost twice as likely (42.2%) to report fair or poor oral health compared with those who visited a private dentist (24.6%).
- People with no dental insurance were almost twice as likely (21.9%) to report fair or poor oral health compared with those with insurance (11.8%).

Discussion

Dentate residents of NSW were equally as likely as other Australians to report that their oral health was 'fair' or 'poor'. Reporting fair or poor oral health was associated with living in a low socioeconomic postcode, being a government health cardholder, having last visited a public dental service and not having dental insurance.

Table 28: Percentage of people rating their oral health fair or poor

		Population: dentate people Age (years)			
		All ages	15–34	35–54	≥55
All people	Per cent of people	16.9	14.0	19.1	18.0
	95% CI ^(a)	15.1–18.8	11.3–17.1	16.2–22.2	15.4–20.8
Sex					
Males	% of people	18.0	14.5	20.9	18.8
	95% CI	15.6–20.6	10.6–19.6	16.8–25.6	15.2–23.0
Females	% of people	15.8	13.4	17.2	17.2
	95% CI	13.8–18.0	10.4–17.1	14.3–20.6	14.2–20.6
Residential location					
Capital city	% of people	16.0	14.2	17.3	17.0
	95% CI	14.0–18.3	10.9–18.2	14.2–21.0	13.5–21.2
Other places	% of people	18.4	13.5	22.1	19.4
	95% CI	15.2–22.2	9.4–19.0	16.9–28.3	16.1–23.1
Postcode socioeconomic status					
Lowest	% of people	21.1	15.1	25.6	22.2
	95% CI	17.9–24.6	11.0–20.4	19.8–32.3	18.4–26.5
Middle	% of people	15.7	12.9	17.0	18.2
	95% CI	12.4–19.6	8.4–19.3	13.1–21.8	13.3–24.4
Highest	% of people	14.4	14.2	15.2	13.6
	95% CI	12.3–16.8	10.1–19.5	11.4–19.9	10.0–18.2
Government health card					
Health care card or pensioner concession card	% of people	29.6	30.9	40.1	24.5
	95% CI	25.5–34.1	22.2–41.2	30.9–50.1	20.3–29.2
Neither card	% of people	13.2	11.0	15.6	12.8
	95% CI	11.7–14.9	8.6–14.0	13.2–18.4	10.4–15.7
Place of last dental visit					
Cardholder/Public	% of people	42.2	36.7	57.2	35.2
	95% CI	34.7–50.2	22.9–53.2	43.3–70.0	26.3–45.2
Cardholder/Non-public	% of people	24.6	26.2	30.2	22.3
	95% CI	20.8–28.9	16.7–38.6	21.3–41.0	17.9–27.4
Non-cardholder/Non-public	% of people	13.2	11.0	15.6	12.8
	95% CI	11.7–14.9	8.6–14.0	13.2–18.4	10.4–15.7
Dental insurance					
Insured	% of people	11.8	10.0	11.7	14.3
	95% CI	10.1–13.7	6.7–14.5	9.5–14.5	11.5–17.8
Uninsured	% of people	21.9	17.5	28.3	20.8
	95% CI	19.4–24.7	13.8–22.0	24.0–33.1	17.2–24.8

(a) 95% CI = 95% confidence interval for estimated percentage.

Percentage of people experiencing toothache

Toothache is caused when the nerve root of a tooth is irritated. It is most commonly caused by infection, decay, injury or loss of a tooth. However, pain sometimes originates from other areas, most commonly the jaw joint and the ear, and radiates to the jaw, thus appearing to be tooth pain.

In NSAOH, experience of toothache was assessed in the interview by asking dentate people 'During the last 12 months how often have you had toothache? Was it: very often, often, sometimes, hardly ever, never during the last 12 months, don't know?'. People who answered 'very often', 'often' or 'sometimes' were classified as having experienced toothache. They represented 16.2% of the dentate NSW population aged 15 years or more (Table 29), which was slightly higher than the national estimate of 15.1% (Slade et al. 2007). The difference was not statistically significant.

Key findings

- The experience of toothache decreased with age, from 21.2% in the 15–34-year-olds to 10.3 % in those aged 55 years or more.
- Experience of toothache was higher in government health cardholders (22.6%) than non-government health cardholders (14.4%). There was a three-fold difference in the 35–44 years age group compared with a 1.5-fold difference in the other age groups.
- Within the population of government health cardholders, those who last visited a public dental clinic were almost three times as likely (42.9%) to report experience of toothache compared with those who visited a private dentist (14.6%). The latter group had the same experience of toothache as non-government health cardholders.

Discussion

Residents of NSW were equally as likely as the rest of the Australian population to experience toothache. Experience of toothache was associated with being young and having last visited a public dental clinic.

Table 29: Percentage of people experiencing toothache

		Population: dentate people Age (years)			
		All ages	15–34	35–54	≥55
All people	Per cent of people	16.2	21.2	15.6	10.3
	<i>95% CI^(a)</i>	<i>14.5–18.2</i>	<i>18.0–24.9</i>	<i>13.3–18.2</i>	<i>8.4–12.6</i>
Sex					
Males	% of people	15.0	17.4	15.8	10.5
	<i>95% CI</i>	<i>12.8–17.4</i>	<i>13.2–22.6</i>	<i>12.4–19.9</i>	<i>7.9–13.7</i>
Females	% of people	17.5	25.1	15.4	10.2
	<i>95% CI</i>	<i>15.4–19.8</i>	<i>21.5–29.2</i>	<i>13.0–18.1</i>	<i>7.6–13.5</i>
Residential location					
Capital city	% of people	16.2	21.1	15.5	9.7
	<i>95% CI</i>	<i>13.9–18.7</i>	<i>16.9–25.9</i>	<i>12.7–18.7</i>	<i>7.5–12.5</i>
Other places	% of people	16.3	21.6	15.8	11.2
	<i>95% CI</i>	<i>13.8–19.2</i>	<i>17.1–27.1</i>	<i>12.1–20.3</i>	<i>8.2–15.2</i>
Postcode socioeconomic status					
Lowest	% of people	18.7	24.0	18.2	13.2
	<i>95% CI</i>	<i>15.3–22.6</i>	<i>18.8–30.0</i>	<i>13.7–23.6</i>	<i>9.4–18.2</i>
Middle	% of people	16.3	20.5	16.1	9.5
	<i>95% CI</i>	<i>13.5–19.5</i>	<i>15.4–26.8</i>	<i>12.3–20.7</i>	<i>7.0–12.9</i>
Highest	% of people	14.0	19.8	12.7	8.2
	<i>95% CI</i>	<i>11.5–16.9</i>	<i>14.2–26.8</i>	<i>9.9–16.2</i>	<i>5.8–11.6</i>
Government health card					
Health care card or pensioner concession card	% of people	22.6	30.3	37.0	12.8
	<i>95% CI</i>	<i>19.1–26.6</i>	<i>22.1–39.9</i>	<i>29.3–45.5</i>	<i>9.6–16.9</i>
Neither card	% of people	14.4	19.6	12.1	8.4
	<i>95% CI</i>	<i>12.7–16.2</i>	<i>16.3–23.5</i>	<i>10.0–14.6</i>	<i>6.3–11.0</i>
Place of last dental visit					
Cardholder/Public	% of people	42.9	44.0	55.7	29.7
	<i>95% CI</i>	<i>35.2–51.0</i>	<i>29.3–59.8</i>	<i>41.7–68.8</i>	<i>20.6–40.8</i>
Cardholder/Non-public	% of people	14.6	19.1	26.2	9.4
	<i>95% CI</i>	<i>11.6–18.2</i>	<i>12.0–29.1</i>	<i>18.6–35.6</i>	<i>6.8–12.8</i>
Non-cardholder/Non-public	% of people	14.4	19.6	12.1	8.4
	<i>95% CI</i>	<i>12.7–16.2</i>	<i>16.3–23.5</i>	<i>10.0–14.6</i>	<i>6.3–11.0</i>
Dental insurance					
Insured	% of people	14.0	22.5	10.9	8.0
	<i>95% CI</i>	<i>11.8–16.4</i>	<i>17.9–27.8</i>	<i>8.6–13.9</i>	<i>5.8–10.9</i>
Uninsured	% of people	18.4	20.4	21.5	12.3
	<i>95% CI</i>	<i>16.2–20.9</i>	<i>16.2–25.4</i>	<i>18.3–25.1</i>	<i>9.7–15.6</i>

(a) 95% CI = 95% confidence interval for estimated percentage.

Percentage of people experiencing orofacial pain

Orofacial pain can be debilitating and indicates temporomandibular joint dysfunction.

In NSAOH, orofacial pain was assessed in the interview by asking people 'During the last month, have you had pain in the face, jaw, temple, in front of the ear or in the ear?'. People who answered 'yes' were classified as having orofacial pain. They represented 23.3% of the NSW population aged 15 years or more (Table 30), which was slightly higher than the national estimate of 22.6 (Slade et al. 2007). The difference was not statistically significant.

Key findings

- The experience of orofacial pain decreased with age, from 25.4% in the 15–34-year olds to 19.0 % in those aged 55 years or more.
- Females were more likely to report that they had orofacial pain (27.3%) than males (19.1%).

Discussion

Residents of NSW equally as likely to experience orofacial pain as the rest of the Australian population. Experience of orofacial pain was associated with being young and being female.

Table 30: Percentage of people experiencing orofacial pain

		Population: all people Age (years)			
		All ages	15–34	35–54	≥55
All people	Per cent of people	23.3	25.4	24.8	19.0
	<i>95% CI^(a)</i>	21.5–25.1	22.1–29.0	22.3–27.5	16.8–21.5
Sex					
Males	% of people	19.1	19.6	21.9	15.0
	<i>95% CI</i>	16.7–21.7	15.2–25.0	18.2–26.2	11.9–18.8
Females	% of people	27.3	31.4	27.6	22.6
	<i>95% CI</i>	25.2–29.5	26.9–36.1	24.4–31.1	19.9–25.6
Residential location					
Capital city	% of people	22.9	25.1	23.4	19.3
	<i>95% CI</i>	20.7–25.3	21.2–29.4	20.2–27.0	16.0–23.0
Other places	% of people	23.9	26.1	27.1	18.7
	<i>95% CI</i>	21.2–26.8	20.3–32.8	23.4–31.2	16.0–21.8
Postcode socioeconomic status					
Lowest	% of people	25.4	29.1	27.6	20.0
	<i>95% CI</i>	22.8–28.3	23.5–35.5	24.0–31.4	17.1–23.2
Middle	% of people	21.5	21.7	22.2	20.5
	<i>95% CI</i>	18.8–24.6	16.9–27.4	18.3–26.6	15.9–26.1
Highest	% of people	22.9	26.3	24.8	16.5
	<i>95% CI</i>	19.7–26.5	20.8–32.8	20.0–30.4	13.0–20.8
Government health card					
Health care card or pensioner concession card	% of people	24.7	29.5	32.3	20.4
	<i>95% CI</i>	21.7–28.0	20.6–40.3	25.3–40.3	17.6–23.7
Neither card	% of people	22.9	24.9	23.5	17.8
	<i>95% CI</i>	20.8–25.0	21.4–28.8	20.7–26.4	14.5–21.5
Place of last dental visit					
Cardholder/Public	% of people	28.8	31.1	33.1	23.6
	<i>95% CI</i>	22.8–35.5	19.1–46.3	21.8–46.8	17.4–31.2
Cardholder/Non-public	% of people	23.2	28.2	31.9	19.7
	<i>95% CI</i>	19.9–26.7	18.1–41.1	23.7–41.4	16.6–23.3
Non-cardholder/Non-public	% of people	22.9	24.9	23.5	17.8
	<i>95% CI</i>	20.8–25.0	21.4–28.8	20.7–26.4	14.5–21.5
Dental insurance					
Insured	% of people	21.8	23.5	22.9	18.1
	<i>95% CI</i>	19.5–24.3	18.9–28.8	19.6–26.5	15.1–21.6
Uninsured	% of people	24.9	27.9	27.3	19.7
	<i>95% CI</i>	22.5–27.5	23.2–33.1	23.4–31.5	17.0–22.8

(a) 95% CI = 95% confidence interval for estimated percentage.

Perceived need for dentures

In NSAOH people were asked at the time of the interview, 'Currently, which of the following dental treatments do you think you need to have?'. The possible responses varied for dentate and edentulous people. All people were asked if they felt they needed dentures.

In NSW 7.1% of people thought they needed dentures (Table 31), which was very similar to the national estimate of 7.2% (Slade et al. 2007).

Key findings

- The percentage of adults who thought they needed dentures was strongly age-related, increasing from 1.3% in the 15–34 years age group to 4.6% among 35–54-year-olds and 16.5% in the 55 years or more age group.
- People living outside a capital city (10.0%) were nearly twice as likely to need a denture as those living in a capital city (5.4%).
- The percentage who thought they needed a denture decreased from people living in postcodes of low socioeconomic status (10.9%) to those living in postcodes of high socioeconomic status (3.4%).
- The need for a denture was five times higher among government health cardholders (17.9%) compared with non-government health cardholders (3.4%).
- Those who had a government health card who last visited a public clinic reported the highest percentage (25.0%). The percentage was lower among people who had a government health card who last visited a private dentist (15.2%) and lowest among those who did not have a government health card and last visited a private dentist (3.4%).
- The insured were less likely to report a need for dentures (3.5%), while among those without dental insurance the need was three times higher (10.4%).
- The age-relatedness of the need for dentures was evident within subgroups of adults formed by socioeconomic characteristics. For instance, among people without dental insurance, the percentage rose from 1.6% in the 15–34 years age group to 8.6% among 35–54-year-olds and 21.4% in the 55 years old or more age group.
- Many of these differences by socioeconomic characteristic persisted when subgroups of adults within the oldest age group (55 years or more) were compared.

Discussion

Perceived need for dentures was low. It is related to the observed pattern for complete tooth loss and numbers of missing teeth. However, the level of need for dentures was considerably lower than the percentage of people with either complete tooth loss or reasonable numbers of missing teeth. The relationship between perceived need and professional judgement of the need for dentures is complex, but people generally express a lower need than is assessed by dentists.

Table 31: Percentage of people who need dentures

		Population: all people Age (years)			
		All ages	15–34	35–54	≥55
All people	Per cent of people	7.1	1.3	4.6	16.5
	<i>95% CI^(a)</i>	<i>6.2–8.1</i>	<i>0.6–3.2</i>	<i>3.4–6.2</i>	<i>14.2–19.1</i>
Sex					
Males	% of people	6.9	1.7	4.9	15.7
	<i>95% CI</i>	<i>5.6–8.5</i>	<i>0.5–6.0</i>	<i>3.1–7.8</i>	<i>12.7–19.2</i>
Females	% of people	7.2	1.0	4.3	17.2
	<i>95% CI</i>	<i>6.2–8.5</i>	<i>0.4–2.3</i>	<i>2.9–6.2</i>	<i>14.6–20.2</i>
Residential location					
Capital city	% of people	5.4	1.1	4.0	12.8
	<i>95% CI</i>	<i>4.4–6.6</i>	<i>0.5–2.6</i>	<i>2.8–5.9</i>	<i>10.0–16.3</i>
Other places	% of people	10.0	1.8	5.6	21.4
	<i>95% CI</i>	<i>8.3–12.0</i>	<i>0.3–9.2</i>	<i>3.4–8.9</i>	<i>17.9–25.5</i>
Postcode socioeconomic status					
Lowest	% of people	10.9	1.9	7.0	22.8
	<i>95% CI</i>	<i>9.0–13.1</i>	<i>0.3–10.6</i>	<i>4.6–10.4</i>	<i>19.2–26.8</i>
Middle	% of people	7.1	1.1	4.8	18.3
	<i>95% CI</i>	<i>5.8–8.7</i>	<i>0.4–3.6</i>	<i>2.9–8.0</i>	<i>14.5–22.9</i>
Highest	% of people	3.4	1.1	2.2	7.6
	<i>95% CI</i>	<i>2.4–4.7</i>	<i>0.3–3.4</i>	<i>1.2–4.1</i>	<i>5.0–11.4</i>
Government health card					
Health care card or pensioner concession card	% of people	17.9	2.7	16.2	23.9
	<i>95% CI</i>	<i>15.5–20.6</i>	<i>1.0–7.1</i>	<i>11.3–22.7</i>	<i>20.6–27.6</i>
Neither card	% of people	3.4	1.1	2.6	9.3
	<i>95% CI</i>	<i>2.7–4.4</i>	<i>0.3–3.6</i>	<i>1.8–3.8</i>	<i>7.2–11.9</i>
Place of last dental visit					
Cardholder/Public	% of people	25.0	2.9	27.9	42.0
	<i>95% CI</i>	<i>19.3–31.7</i>	<i>0.8–10.0</i>	<i>16.5–43.1</i>	<i>32.9–51.7</i>
Cardholder/Non-public	% of people	15.2	2.5	8.9	19.9
	<i>95% CI</i>	<i>12.6–18.2</i>	<i>0.6–10.6</i>	<i>4.9–15.6</i>	<i>16.5–23.7</i>
Non-cardholder/Non-public	% of people	3.4	1.1	2.6	9.3
	<i>95% CI</i>	<i>2.7–4.4</i>	<i>0.3–3.6</i>	<i>1.8–3.8</i>	<i>7.2–11.9</i>
Dental insurance					
Insured	% of people	3.5	1.1	1.3	9.5
	<i>95% CI</i>	<i>2.7–4.5</i>	<i>0.4–3.4</i>	<i>0.7–2.6</i>	<i>7.0–12.8</i>
Uninsured	% of people	10.4	1.6	8.6	21.4
	<i>95% CI</i>	<i>8.9–12.1</i>	<i>0.4–5.3</i>	<i>6.2–11.8</i>	<i>18.6–24.6</i>

(a) 95% CI = 95% confidence interval for estimated percentage.

Perceived need for dental extraction or filling

Dentate adults were asked about additional dental services, including extractions or fillings that they might need. The responses to the options 'Any extractions' or 'Any fillings' have been combined so that the response indicates a perceived dental problem for which one or other of these two aspects of routine dental care was thought to be required, most likely as a sequelae for dental caries. Which of these two dental services were provided would be determined by a process of negotiation between patient and provider that would be influenced by the specific circumstances.

In NSW, 31.1% of dentate adults perceived a need for an extraction or filling (Table 32), which was similar to the national estimate of 32.9% (Slade et al. 2007).

Key findings

- The percentage of dentate adults who thought they needed extractions or fillings was similar across all three age groups, varying from 34.0% among people aged 35–54 years to 27.3% among people aged 55 years or more.
- There were no significant differences by sex or residential location.
- The percentage of dentate adults who thought they needed an extraction or filling was similar for people living in postcodes with low or middle socioeconomic status (33.5% and 33.2% respectively) but was lower among those living in postcodes of high socioeconomic status (26.8%).
- The need for an extraction or filling was higher among people who were government health cardholders (39.4%) compared with those who did not have a government health card (28.7%).
- Among government health cardholders, those who last visited a public clinic had the highest percentage of a need for an extraction or filling (53.1%). The percentage was significantly lower among cardholders who last visited a private dentist (34.1%) and those adults who did not have a government health card who last visited a private dentist (28.7%).
- People who were uninsured had a higher perceived need for extractions (35.9%) than those who were insured (26.3%).
- The lack of an age-related pattern of need for an extraction or filling was repeated within subgroups of dentate adults formed by socioeconomic characteristics. Occasionally the trend for a lower percentage in need among those aged 55 years or more reached significance. For instance, among those with government health cards, this age group had a significantly lower percentage than the other age groups.

Discussion

About one-third of dentate adults perceived a need for an extraction or filling. This showed no significant variation by age and few socioeconomic characteristic variations. The percentage was higher among people living in middle or lower socioeconomic status postcodes, government health cardholders who last visited a public dental clinic and the uninsured.

Table 32: Percentage of people who need an extraction or filling

		Population: dentate people Age (years)			
		All ages	15–34	35–54	≥55
All people	Per cent of people	31.1	31.0	34.0	27.3
	<i>95% CI^(a)</i>	<i>29.3–32.9</i>	<i>27.9–34.3</i>	<i>31.1–37.0</i>	<i>24.9–29.9</i>
Sex					
Males	% of people	31.8	30.0	34.5	30.7
	<i>95% CI</i>	<i>29.1–34.7</i>	<i>25.1–35.4</i>	<i>30.2–39.1</i>	<i>26.5–35.2</i>
Females	% of people	30.4	32.0	33.4	24.0
	<i>95% CI</i>	<i>27.7–33.1</i>	<i>27.4–37.1</i>	<i>29.7–37.4</i>	<i>21.0–27.4</i>
Residential location					
Capital city	% of people	29.6	30.2	31.5	25.9
	<i>95% CI</i>	<i>27.4–31.9</i>	<i>26.4–34.4</i>	<i>28.0–35.2</i>	<i>22.6–29.5</i>
Other places	% of people	33.8	32.6	38.4	29.4
	<i>95% CI</i>	<i>30.9–36.8</i>	<i>27.6–38.2</i>	<i>33.6–43.5</i>	<i>26.1–33.0</i>
Postcode socioeconomic status					
Lowest	% of people	33.5	30.6	37.9	31.2
	<i>95% CI</i>	<i>30.9–36.1</i>	<i>25.5–36.2</i>	<i>33.2–42.9</i>	<i>27.6–35.0</i>
Middle	% of people	33.2	31.7	38.5	27.7
	<i>95% CI</i>	<i>30.4–36.2</i>	<i>26.7–37.1</i>	<i>34.3–42.9</i>	<i>23.2–32.8</i>
Highest	% of people	26.8	30.6	25.8	23.3
	<i>95% CI</i>	<i>23.6–30.3</i>	<i>24.9–36.9</i>	<i>21.0–31.2</i>	<i>19.6–27.3</i>
Government health card					
Health care card or pensioner concession card	% of people	39.4	46.4	53.2	30.3
	<i>95% CI</i>	<i>35.6–43.4</i>	<i>37.1–55.9</i>	<i>44.0–62.1</i>	<i>26.8–33.9</i>
Neither card	% of people	28.7	28.3	30.9	25.0
	<i>95% CI</i>	<i>26.6–30.9</i>	<i>24.8–32.1</i>	<i>28.2–33.7</i>	<i>21.7–28.7</i>
Place of last dental visit					
Cardholder/Public	% of people	53.1	53.5	64.6	41.7
	<i>95% CI</i>	<i>44.6–61.4</i>	<i>37.7–68.7</i>	<i>51.3–75.9</i>	<i>31.7–52.5</i>
Cardholder/Non-public	% of people	34.1	40.6	46.5	27.9
	<i>95% CI</i>	<i>30.4–37.9</i>	<i>30.2–51.9</i>	<i>35.7–57.6</i>	<i>24.2–32.0</i>
Non-cardholder/Non-public	% of people	28.7	28.3	30.9	25.0
	<i>95% CI</i>	<i>26.6–30.9</i>	<i>24.8–32.1</i>	<i>28.2–33.7</i>	<i>21.7–28.7</i>
Dental insurance					
Insured	% of people	26.3	28.5	27.1	22.3
	<i>95% CI</i>	<i>24.0–28.9</i>	<i>23.8–33.8</i>	<i>23.8–30.6</i>	<i>19.0–26.0</i>
Uninsured	% of people	35.9	33.5	42.6	31.6
	<i>95% CI</i>	<i>33.4–38.5</i>	<i>29.1–38.3</i>	<i>38.1–47.3</i>	<i>28.2–35.2</i>

(a) 95%CI = 95% confidence interval for estimated percentage.

Perceived need for a dental check-up

Dentate adults were asked about their perceived need for a check-up. This is regarded as an indicator of compliance with the recommendation of dentists to visit regularly when asymptomatic so as to detect disease earlier and receive prompt treatment for any dental problems. A check-up also provides an opportunity for preventive services to be received.

In NSW, 58.0% of dentate adults perceived a need for a check-up (Table 33), which was similar to the national estimate of 59.6% (Slade et al. 2007).

Key findings

- The percentage of dentate adults who thought they needed a check-up was similar across the two younger age groups (63.2 and 60.3%) but lower among people aged 55 years or more (48.0%).
- There were no significant differences among all dentate adults by sex, residential location, postcode socioeconomic status, government health cardholder status or place of last dental visit.
- People who were uninsured were more likely to perceive need for a check-up (62.6%) than those who were insured (53.2%).
- The age-related pattern of perceived need for a check-up was repeated within subgroups of dentate adults formed by most of the socioeconomic characteristics. Only among subgroups formed by place of last visit was the age-related pattern not significantly lower among adults aged 55 years or more.

Discussion

About 6 out of 10 dentate adults perceived a need for a check-up. The percentage was similar for the two younger age groups but significantly lower among people aged 55 years or more. There was little variation by socioeconomic characteristics, which might reflect a confounding of perceived need for a check-up by time since last dental visit. Those with a higher likelihood of compliance with the recommendation of a regular check-up visit may have last visited more recently, and hence not perceive a need for a further check-up at the time of the interview.

Table 33: Percentage of people perceiving a need for check-up

		Population: dentate people			
		Age (years)			
		All ages	15–34	35–54	≥55
All people	Per cent of people	58.0	63.2	60.3	48.0
	<i>95% CI^(a)</i>	<i>56.0–60.0</i>	<i>59.2–66.9</i>	<i>57.3–63.3</i>	<i>44.9–51.2</i>
Sex					
Males	% of people	57.5	59.4	62.7	47.8
	<i>95% CI</i>	<i>54.6–60.4</i>	<i>53.4–65.2</i>	<i>58.1–67.0</i>	<i>43.4–52.2</i>
Females	% of people	58.6	67.0	58.0	48.3
	<i>95% CI</i>	<i>55.8–61.4</i>	<i>62.0–71.6</i>	<i>54.2–61.6</i>	<i>43.7–52.9</i>
Residential location					
Capital city	% of people	57.5	61.4	59.5	48.6
	<i>95% CI</i>	<i>54.9–60.1</i>	<i>56.6–66.0</i>	<i>55.8–63.2</i>	<i>44.3–53.0</i>
Other places	% of people	59.0	66.8	61.7	47.2
	<i>95% CI</i>	<i>55.8–62.0</i>	<i>60.1–72.9</i>	<i>56.3–66.8</i>	<i>42.9–51.5</i>
Postcode socioeconomic status					
Lowest	% of people	59.5	65.6	62.4	49.0
	<i>95% CI</i>	<i>56.3–62.6</i>	<i>58.1–72.3</i>	<i>56.4–68.1</i>	<i>44.2–53.9</i>
Middle	% of people	58.2	63.1	59.7	47.9
	<i>95% CI</i>	<i>54.7–61.6</i>	<i>57.0–68.8</i>	<i>55.0–64.3</i>	<i>42.2–53.7</i>
Highest	% of people	56.5	61.1	59.1	47.2
	<i>95% CI</i>	<i>52.9–60.1</i>	<i>54.0–67.9</i>	<i>53.8–64.1</i>	<i>41.6–52.9</i>
Government health card					
Health care card or pensioner concession card	% of people	60.9	71.1	70.7	51.9
	<i>95% CI</i>	<i>57.4–64.4</i>	<i>61.7–79.0</i>	<i>63.2–77.3</i>	<i>47.7–56.2</i>
Neither card	% of people	57.2	61.8	58.7	45.0
	<i>95% CI</i>	<i>54.9–59.5</i>	<i>57.5–66.0</i>	<i>55.5–61.8</i>	<i>40.8–49.2</i>
Place of last dental visit					
Cardholder/Public	% of people	67.3	77.9	70.9	50.6
	<i>95% CI</i>	<i>60.9–73.0</i>	<i>63.8–87.6</i>	<i>57.9–81.2</i>	<i>40.2–60.9</i>
Cardholder/Non-public	% of people	58.5	65.7	70.6	52.2
	<i>95% CI</i>	<i>54.3–62.5</i>	<i>51.9–77.2</i>	<i>61.7–78.2</i>	<i>47.8–56.6</i>
Non-cardholder/Non-public	% of people	57.2	61.8	58.7	45.0
	<i>95% CI</i>	<i>54.9–59.5</i>	<i>57.5–66.0</i>	<i>55.5–61.8</i>	<i>40.8–49.2</i>
Dental insurance					
Insured	% of people	53.2	58.5	54.7	43.8
	<i>95% CI</i>	<i>50.1–56.2</i>	<i>52.0–64.7</i>	<i>50.7–58.6</i>	<i>39.7–48.0</i>
Uninsured	% of people	62.6	67.3	67.0	51.4
	<i>95% CI</i>	<i>60.1–65.1</i>	<i>62.5–71.8</i>	<i>62.9–70.9</i>	<i>47.2–55.7</i>

(a) 95% CI = 95% confidence interval for estimated percentage.

Perceived urgency of dental treatment needs

Dentate adults who perceived a need for an extraction or filling were asked about the urgency of needed dental treatment. Dental problems vary from truly urgent problems like dental trauma, swelling in or around the jaws, or bleeding (usually as a complication of dental treatment) through situations where treatment is highly desirable in a short period of time (usually associated with pain) to problems that can reasonably wait to be treated. In NSAOH, dentate adults who perceived a need for an extraction or filling were asked 'How soon do you think you need this dental treatment?' at the time of the interview. The possible responses included a wide range of time periods. These have been collapsed to perceiving a need for treatment either within 3 months or longer than 3 months.

In NSW, 71.6% of dentate adults needing an extraction or filling perceived a need for dental treatment within 3 months (Table 34), which was similar to the national estimate of 69.3% (Slade et al. 2007).

Key findings

- The percentage of dentate adults needing an extraction or filling who thought they needed dental treatment within 3 months showed little variation by age group. The percentage varied only from 70.3% to 72.9% across the three age groups.
- There were no significant differences among subgroups formed by any social characteristic.

Discussion

About 7 out of 10 dentate adults who needed an extraction or filling perceived a need for dental treatment within 3 months. The percentage was similar for the three age groups. Perceived need for dental treatment showed little variation by socioeconomic characteristics, which might reflect a confounding of perceived need for dental treatment within 3 months by time since last dental visit.

Table 34: Percentage of people perceiving a need for treatment within 3 months

		Population: dentate people who need an extraction or filling			
		Age (years)			
		All ages	15–34	35–54	≥55
All people	Per cent of people	71.6	72.2	70.3	72.9
	<i>95% CI^(a)</i>	<i>67.7–75.2</i>	<i>64.2–78.9</i>	<i>64.9–75.1</i>	<i>66.3–78.5</i>
Sex					
Males	% of people	70.1	73.1	68.6	68.1
	<i>95% CI</i>	<i>64.6–75.2</i>	<i>59.8–83.3</i>	<i>60.6–75.6</i>	<i>59.6–75.6</i>
Females	% of people	73.1	71.2	71.9	78.9
	<i>95% CI</i>	<i>68.3–77.4</i>	<i>62.9–78.3</i>	<i>65.4–77.6</i>	<i>70.1–85.6</i>
Residential location					
Capital city	% of people	70.6	70.2	69.2	74.0
	<i>95% CI</i>	<i>65.8–75.0</i>	<i>60.7–78.3</i>	<i>62.4–75.3</i>	<i>64.0–82.0</i>
Other places	% of people	73.1	76.0	71.9	71.3
	<i>95% CI</i>	<i>66.3–78.9</i>	<i>60.8–86.6</i>	<i>62.9–79.4</i>	<i>63.2–78.3</i>
Postcode socioeconomic status					
Lowest	% of people	70.0	73.0	70.9	65.1
	<i>95% CI</i>	<i>63.3–76.0</i>	<i>59.3–83.3</i>	<i>61.8–78.6</i>	<i>56.6–72.6</i>
Middle	% of people	72.2	74.3	67.8	78.1
	<i>95% CI</i>	<i>65.9–77.7</i>	<i>61.4–83.9</i>	<i>58.5–75.8</i>	<i>65.9–86.7</i>
Highest	% of people	72.6	69.1	73.2	77.7
	<i>95% CI</i>	<i>64.6–79.3</i>	<i>53.5–81.3</i>	<i>63.2–81.4</i>	<i>63.7–87.4</i>
Government health card					
Health care card or pensioner concession card	% of people	73.4	72.4	75.5	72.6
	<i>95% CI</i>	<i>67.1–78.9</i>	<i>59.0–82.7</i>	<i>64.4–84.0</i>	<i>64.3–79.6</i>
Neither card	% of people	71.1	72.6	68.9	73.1
	<i>95% CI</i>	<i>66.4–75.3</i>	<i>63.7–80.0</i>	<i>62.8–74.4</i>	<i>63.1–81.1</i>
Place of last dental visit					
Cardholder/Public	% of people	83.2	83.0	77.8	91.6
	<i>95% CI</i>	<i>72.7–90.2</i>	<i>60.5–94.0</i>	<i>59.4–89.4</i>	<i>79.2–96.9</i>
Cardholder/Non-public	% of people	67.7	61.6	73.8	67.4
	<i>95% CI</i>	<i>59.7–74.8</i>	<i>42.9–77.4</i>	<i>59.7–84.2</i>	<i>57.6–76.0</i>
Non-cardholder/Non-public	% of people	71.1	72.6	68.9	73.1
	<i>95% CI</i>	<i>66.4–75.3</i>	<i>63.7–80.0</i>	<i>62.8–74.4</i>	<i>63.1–81.1</i>
Dental insurance					
Insured	% of people	73.7	77.4	68.5	78.1
	<i>95% CI</i>	<i>68.5–78.4</i>	<i>67.3–85.1</i>	<i>60.6–75.5</i>	<i>65.4–87.1</i>
Uninsured	% of people	69.8	68.3	71.6	69.4
	<i>95% CI</i>	<i>64.7–74.5</i>	<i>57.4–77.4</i>	<i>65.0–77.5</i>	<i>61.2–76.5</i>

(a) 95% CI = 95% confidence interval for estimated percentage.

Age-standardised comparison between government health cardholders and non-health cardholders

Findings from 29 of the preceding tables are summarised in Table 35, to compare oral health indicators between people with a government health card and non-cardholders. Percentages and means for the two groups are age-standardised, a statistical procedure that aims to remove any effects of age that might account for differences between the two groups in each oral health indicator. As noted in Table 4, a smaller percentage of people in the youngest age group had a health care card or pensioner concession card than in the oldest age group. Age standardisation seeks to compensate for that difference in age distribution, so that differences in any single indicator between the two groups are not confounded by age.

- For most outcomes reported in Table 35, health cardholders had significantly poorer oral health status, oral health care and perceived oral health.
- Exceptions occurred for untreated root decay, average number of DMF teeth per person and four periodontal conditions, where the slightly greater frequency in health cardholders compared with non-cardholders was not statistically significant. Also, the slightly lower percentage with filled teeth among health cardholders compared with non-cardholders was not statistically significant.
- For measures relating to tooth loss, the magnitude of difference in age-standardised estimates between the two groups was noticeably smaller than the difference between the same two groups noted in preceding tables where there was no adjustment for age. For example, health cardholders had a 2.6-fold greater prevalence of complete tooth loss when the comparison was adjusted for age (Table 35), whereas prevalence differed by a factor of 7.6 when all ages were contrasted in Table 5 (15.4% for health cardholders compare with 2.0% for non-cardholders). This degree of attenuation indicates that age was an important confounder of the relationship between health card status and complete tooth loss.
- However, for most other indicators in Table 35, the relative differences in age-standardised results between the two groups were similar in magnitude to preceding tables. This is because there was only a weak association between age and dental attendance, with the consequence that there was little confounding of the difference between the two groups by age.

In summary, the findings in Table 35 confirm that health cardholders are disadvantaged with respect to most indicators of oral health status, oral health care and perceived oral health, and that the disadvantage is not due to the older age profile of health cardholders compared to non-cardholders. Even when age standardisation attenuated the difference between the two groups, as observed for measures relating to tooth loss, the differences tended to remain statistically significant.

Table 35: Age-standardised comparison of health cardholders and non-health cardholders

Variable	Cardholders	Non-cardholders
	Estimate (95%CI)	Estimate (95%CI)
% of people with complete tooth loss	8.6 (6.8–10.4)	3.3 (2.5–4.0)
% of people with fewer than 21 teeth	21.1 (18.4–23.8)	9.1 (7.7–10.5)
% of dentate people who wear denture(s)	21.4 (18.8–24.0)	14.8 (13.0–16.6)
Average number of missing teeth per person	6.5 (5.7–7.2)	4.6 (4.1–5.1)
% of people with untreated coronal decay	40.8 (32.0–49.6)	23.5 (19.2–27.7)
% of people with untreated root decay	12.0 (7.9–16.0)	7.2 (4.9–9.5)
% of people with one or more filled teeth	73.1 (68.3–77.8)	81.3 (77.1–85.5)
Average number of DMF teeth per person	14.1 (13.3–15.0)	12.9 (12.4–13.5)
% of people with moderate or severe periodontitis	32.0 (25.3–38.6)	22.0 (19.1–24.8)
% of people with 4+ mm periodontal pocket depth	22.2 (15.7–28.7)	20.2 (16.8–23.6)
% of people with 4+ mm clinical attachment loss	50.8 (44.4–57.2)	45.0 (40.9–49.0)
% of people with gingival inflammation	25.5 (16.9–34.1)	15.6 (11.6–19.6)
% of people visiting dentist within last 12 months	52.7 (47.9–57.6)	64.1 (61.8–66.3)
% of people who attended a private dental practice at last dental visit	60.1 (55.1–65.1)	86.9 (85.2–88.6)
% of people who paid for their last dental visit	63.3 (57.8–68.8)	100.0 (100.0–100.0)
% of people who usually visit a dental professional at least once a year	42.6 (38.5–46.7)	59.7 (57.2–62.3)
% of people who have a dentist they usually attend	72.1 (67.8–76.3)	84.5 (82.5–86.5)
% of people who usually visit a dentist for a check up	37.5 (33.1–41.9)	62.7 (60.3–65.2)
% of people who avoided or delayed dental care	46.7 (42.5–51.0)	24.9 (23.0–26.8)
% of people who reported that cost had prevented recommended dental treatment	37.1 (31.7–42.5)	16.6 (14.5–18.7)
% of people who would have a lot of difficulty paying a \$100 dental bill	43.0 (38.0–48.0)	12.5 (11.1–14.0)
% of people avoiding foods due to dental problems	35.2 (30.2–40.2)	15.7 (13.8–17.6)
% of people rating their oral health fair or poor	34.0 (29.1–38.9)	13.1 (11.5–14.7)
% of people experiencing toothache	28.8 (24.4–33.1)	13.0 (11.5–14.5)
% of people experiencing orofacial pain	29.2 (24.9–33.5)	22.2 (20.2–24.3)
% of people who need dentures	14.3 (11.7–16.8)	4.3 (3.2–5.4)
% of people who need an extraction or filling	45.3 (40.4–50.3)	28.2 (26.0–30.3)
% of people perceiving a need for a check up	65.4 (61.6–69.3)	54.9 (52.6–57.3)
% of people perceiving a need for treatment within 3 months	72.7 (66.4–79.1)	72.6 (67.9–77.2)

Age-standardised comparison between the dentally insured and the uninsured

Age standardisation has been used in Table 36 to make comparisons between dentally insured and uninsured people in each of the 30 oral health indicators presented in Tables 5–34. These comparisons are based on the same principles noted for Table 35. That is, age standardisation aims to compare insured and uninsured people after adjusting for potential differences in the age distribution between the two groups. In principle, however, there should be little confounding of these effects because there were only small differences in dental insurance coverage among the three age groups (Table 4).

- The results in Table 36 show statistically significantly poorer outcomes for uninsured people in 23 of the 30 indicators. For each of those 23 indicators, statistically significant differences were also observed in the preceding tables. Furthermore, relative differences between the insured and uninsured for each of those 23 indicators were of a similar magnitude whether or not the comparison was age-standardised.
- Conversely, the seven indicators in Table 36 that did not differ to a statistically significant degree between insured and uninsured people were similarly non-significant when contrasted between the two groups in previous tables that did not use age standardisation.

In summary, the findings in Table 36 confirm generally poorer oral health outcomes for uninsured people compared to insured people. Age standardisation did not appreciably alter the relationship between insurance status and any of the indicators, inferring that there was very little confounding of the effects of insurance due to age.

Table 36: Age-standardised comparison of the dentally insured and the uninsured

Variable	Insured	Uninsured
	Estimate (95%CI)	Estimate (95%CI)
% of people with complete tooth loss	2.8 (2.0–3.7)	7.1 (6.1–8.0)
% of people with fewer than 21 teeth	9.1 (7.5–10.8)	16.9 (15.3–18.6)
% of dentate people who wear denture(s)	14.4 (12.4–16.5)	20.2 (18.3–22.0)
Average number of missing teeth per person	4.4 (3.8–4.9)	5.9 (5.4–6.4)
% of people with untreated coronal decay	21.3 (15.7–26.9)	33.6 (28.4–38.8)
% of people with untreated root decay	4.2 (2.4–5.9)	11.3 (9.0–13.5)
% of people with one or more filled teeth	86.0 (81.3–90.7)	72.9 (68.7–77.1)
Average number of DMF teeth per person	13.3 (12.5–14.1)	13.2 (12.6–13.8)
% of people with moderate or severe periodontitis	21.0 (18.0–24.0)	28.6 (24.4–32.8)
% of people with 4+ mm periodontal pocket depth	19.4 (15.8–23.0)	21.1 (16.7–25.4)
% of people with 4+ mm clinical attachment loss	45.3 (41.0–49.5)	47.2 (42.8–51.6)
% of people with gingival inflammation	13.0 (8.1–17.9)	20.0 (15.7–24.2)
% of people visiting dentist within last 12 months	74.0 (71.5–76.4)	49.4 (46.9–51.8)
% of people who attended a private dental practice at last dental visit	87.9 (85.7–90.1)	74.1 (71.3–77.0)
% of people who paid for their last dental visit	99.0 (98.4–99.6)	85.4 (83.2–87.7)
% of people who received government-subsidised dental care in private sector	0.3 (<0–0.6)	2.8 (2.0–3.5)
% of people who usually visit a dental professional at least once a year	70.0 (66.9–73.0)	41.5 (38.8–44.3)
% of people who have a dentist they usually attend	89.9 (88.0–91.8)	72.4 (69.5–75.2)
% of people who usually visit a dentist for a check up	71.8 (68.8–74.8)	42.9 (40.1–45.7)
% of people who avoided or delayed dental care	17.6 (15.6–19.6)	41.9 (39.1–44.6)
% of people who reported that cost had prevented recommended dental treatment	13.3 (10.8–15.8)	29.3 (25.9–32.6)
% of people who would have a lot of difficulty paying a \$100 dental bill	10.7 (8.9–12.5)	26.8 (24.4–29.2)
% of people avoiding foods due to dental problems	14.2 (12.3–16.1)	24.4 (21.7–27.2)
% of people rating their oral health fair or poor	11.7 (9.8–13.6)	22.2 (19.7–24.7)
% of people experiencing toothache	14.0 (11.7–16.4)	18.4 (16.2–20.7)
% of people experiencing orofacial pain	21.5 (19.1–24.0)	25.5 (23.1–28.0)
% of people who need dentures	3.6 (2.7–4.5)	10.1 (8.6–11.6)
% of people who need an extraction or filling	25.6 (23.1–28.1)	36.4 (33.8–38.9)
% of people perceiving a need for a check up	52.5 (49.6–55.5)	62.7 (60.2–65.1)
% of people perceiving a need for treatment within 3 months	75.3 (70.6–80.0)	69.8 (64.9–74.6)

Appendix

Sample counts

Table A.1: Table counts of interviewed people

	Age group (years)			
	All ages	15–34	35–54	≥55
All people	3,671	791	1,390	1,490
Sex				
Males	1,456	314	539	603
Females	2,215	477	851	887
Residential location				
Capital city	2,246	534	878	834
Other places	1,425	257	512	656
Postcode socioeconomic status				
Lowest	1,213	225	430	558
Middle	1,257	300	492	465
Highest	1,201	266	468	467
Government health card				
Blank but applicable	13	7	2	4
Health care card or pensioner concession card	1,122	132	232	758
Neither card	2,536	652	1156	728
Place of last dental visit				
Cardholder/Public	280	56	79	145
Cardholder/Non-public	842	76	153	613
Dental insurance				
Blank but applicable	27	20	4	3
Insured	1,683	333	746	604
Uninsured	1,961	438	640	883

Table A.2: Sample counts of examined people

	Age group (years)			
	All ages	15–34	35–54	≥55
All people	1,113	176	435	502
Sex				
Males	449	52	167	230
Females	664	124	268	272
Residential location				
Capital city	617	100	256	261
Other places	496	76	179	241
Postcode socioeconomic status				
Lowest	389	56	145	188
Middle	377	61	147	169
Highest	347	59	143	145
Government health card				
Blank but applicable	1	1	0	0
Cardholder	385	44	91	250
Non-cardholder	727	131	344	252
Place of last dental visit				
Cardholder/Public	108	18	31	59
Cardholder/Non-public	277	26	60	191
Dental insurance				
Blank but applicable	7	6	0	1
Insured	491	69	204	218
Uninsured	615	101	231	283

Glossary

95% confidence interval Defines the uncertainty around an estimated value – there is a 95% probability that the true value falls within the range of the upper and lower limits.

Attachment loss The distance in millimetres measured from the edge of the enamel of a tooth to the gum tissue that is adherent to its root.

Calibration A procedure to promote standardisation between examiners performing the oral examinations.

Canine One of four ‘eye teeth’ positioned next to the incisors and used for tearing food.

Capital city The administrative seat of government of each of Australia’s six states and two territories – each capital city also represents the most populous location of its respective state or territory.

Cemento-enamel junction Point on a tooth surface where the tooth crown joins the tooth root.

Census The Census of Population and Housing conducted every 5 years by the Australian Bureau of Statistics.

Complete tooth loss Loss of all natural teeth (also referred to as edentulism).

Coronal Pertaining to the crown of a tooth.

Crown The portion of tooth covered by white enamel that usually is visible in the mouth.

Dental attendance Behaviour related to the use of dental services.

Dental caries The process in which tooth structure is destroyed by acid produced by bacteria in the mouth – see dental decay.

Dental caries experience The cumulative effect of the caries process through a person’s lifetime, manifesting as teeth that are decayed, missing or filled.

Dental decay Cavity resulting from dental caries.

Dental insurance Dental care is not covered under Australia’s universal public health insurance vehicle, Medicare, and consequently people seeking cover can elect to carry private dental insurance.

Dentate Having one or more natural teeth.

Dentition The set of teeth – a complete dentition comprises 32 adult teeth.

Denture A removable dental prosthesis that substitutes for missing natural teeth and adjacent tissues.

DMFT An index of dental caries experience measured by counting the number of decayed (D), missing (M), and filled (F) teeth (T).

Edentulous A state of complete loss of all natural teeth.

Enamel Hard white mineralised tissue covering the crown of a tooth.

Epidemiology The study of the distribution and causes of health and disease in populations.

Examination protocol Methods and guidelines for conducting standardised oral examinations in a survey.

Extraction Removal of a natural tooth.

Generation A group of people born during a defined period of time (also referred to as a birth cohort).

Gingiva Gum tissue.

Gingivitis Redness, swelling or bleeding of the gums caused by inflammation.

Government health card A concession card issued by the Australian Government that entitles the holder to services including public dental care.

Incisor One of eight front teeth used during eating for cutting food.

Index of Relative Socioeconomic Advantage/Disadvantage (IRSAD) One of four indices measuring area-level disadvantage derived by the Australian Bureau of Statistics – the IRSAD is derived from attributes such as low income, low educational attainment, high unemployment and jobs in relatively unskilled occupations.

Indigenous identity A person who states that they are of Aboriginal and/or Torres Strait Islander descent is an Indigenous Australian.

Mean The arithmetic average of a set of values.

Molar One of 12 back teeth used in grinding food.

Natural teeth Refers to a person's own teeth as opposed to artificial teeth.

Orofacial pain Pain located in the face, jaw, temple, in front of the ear or in the ear.

Participation rate The proportion of people from whom survey information is collected from among the total number of people selected as intended study participants.

Periodontal disease Disease of the gums and other tissues that attach to and anchor teeth to the jaws.

Periodontal pocket A space below the gum line that exists between the root of a tooth and the gum surrounding that tooth.

Periodontitis Disease of the gums caused by bacteria, characterised by swelling and bleeding of the gums and loss of tissue that attaches the tooth to the jaw.

Permanent teeth Adult teeth (secondary teeth).

Plaque A film composed of bacteria and food debris that adheres to the tooth surface.

Prevalence The proportion of people with a defined disease within a defined population.

Probing pocket depth The measured depth of the periodontal pocket.

Recorder A person, usually a dental assistant, who recorded the results of an oral examination onto a laptop computer.

Relative difference The difference between two values calculated as a ratio of one value divided by another.

Restoration A filling to repair a tooth damaged by decay or injury.

Root That part of the tooth below the crown which is anchored to the jaw.

Root surface The surface of the root of a tooth.

Socioeconomic Indices for Areas (SEIFA) A set of four indices derived by the Australian Bureau of Statistics from population census data to measure aspects of socioeconomic position for geographic areas.

Socioeconomic position Descriptive term for a position in society and usually measured by attributes such as income, education, occupation or characteristics of residential area.

State/territory Geographic regions of Australia – the nation has six states and two territories.

Statistical significance An indication from a statistical test that an observed association is unlikely (usually less than 5% probability) to be due to chance created when a random sample of people is selected from a population.

Trend The general direction in which change over time is observed.

Unerupted tooth A tooth that has failed to emerge through the gums into the mouth.

Weights Numbers applied to groups of study participants to correct for differences in probability of selection and in participation.

Wisdom tooth One of four molars, each positioned at the back of the mouth.

References

- AHMAC (Australian Health Ministers' Advisory Council) 2001. Steering Committee for National Planning for Oral Health. Oral health of Australians: National planning for oral health improvement. Adelaide: SA Department of Human Services.
- AIHW (Australian Institute of Health and Welfare) 2000. Australia's health 2000: The seventh biennial health report of the Australian Institute of Health and Welfare. Canberra: AIHW.
- AIHW 2007. Health expenditure Australia 2005–06. Health and Welfare Expenditure Series no. 30. Cat. no. HWE 37. Canberra: AIHW.
- Barnard PD 1993. National Oral Health Survey Australia 1987–1988. Canberra: Australian Government Publishing Service.
- Benyamini Y, Leventhal H & Leventhal EA 2004. Self-rated oral health as an independent predictor of self-rated general health, self-esteem and life satisfaction. *Social Science & Medicine* 59(5):1109–16.
- Bergman JD, Wright FA & Hammond RH 1991. The oral health of the elderly in Melbourne. *Australian Dental Journal* 36(4):280–5.
- Brennan DS & Spencer AJ 2004. Changes in caries experience among Australian public dental patients between 1995/96 and 2001/02. *Australian and New Zealand Journal of Public Health* 28(6):542–8.
- Brennan DS, Spencer AJ & Roberts-Thomson KF 2007. Caries experience among 45–54-year-olds in Adelaide, South Australia. *Australian Dental Journal* 52(2):122–7.
- Brennan DS, Spencer AJ & Slade GD 2000. Caries experience among publicly-funded dental patients in Australia, 1995–96: type of care and geographic location. *Australian Dental Journal* 45(1):37–45.
- Brennan DS, Spencer AJ & Slade GD 2001. Prevalence of periodontal conditions among public-funded dental patients in Australia. *Australian Dental Journal* 46(2):114–21.
- Chalmers JM, Carter KD, Fuss JM, Spencer AJ & Hodge CP 2002. Caries experience in existing and new nursing home residents in Adelaide, Australia. *Gerodontology* 19(1):30–40.
- Chalmers JM, Carter KD & Spencer AJ 2002. Caries incidence and increments in community-living older adults with and without dementia. *Gerodontology* 19(2):80–94.
- Chalmers JM, Carter KD & Spencer AJ 2005. Caries incidence and increments in Adelaide nursing home residents. *Special Care Dentistry* 25(2):96–105.
- Chalmers JM, Hodge C, Fuss JM, Spencer AJ & Carter KD 2002. The prevalence and experience of oral diseases in Adelaide nursing home residents. *Australian Dental Journal* 47(2):123–30.
- Coates E, Slade GD, Goss AN & Gorkic E 1996. Oral conditions and their social impact among HIV dental patients. *Australian Dental Journal* 41(1):33–6.
- Coates EA, Brennan D, Logan RM, Goss AN, Scopacasa B, Spencer AJ et al. 2000. Hepatitis C infection and associated oral health problems. *Australian Dental Journal* 45(2):108–14.
- Dawson AS & Smales RJ 1994. Dental health changes in an Australian Defence Force population. *Australian Dental Journal* 39(4):242–6.

- Dillman DA 2000. *Mail and internet surveys: The tailored design method*, 2nd edn. New York: John Wiley Company.
- Elias AC & Sheiham A 1998. The relationship between satisfaction with mouth and number of position of teeth. *Journal of Oral Rehabilitation* 25:649–61.
- Endean C, Roberts-Thomson K & Wooley S 2004. Anangu oral health: The status of the Indigenous population of the Anangu Pitjantjatjara lands. *Australian Journal of Rural Health* 12(3):99–103.
- Hopcraft M & Morgan MV 2003a. Dental caries experience in a young adult military population. *Australian Dental Journal* 48(2):125–9.
- Hopcraft MS & Morgan MV 2003b. Exposure to fluoridated drinking water and dental caries experience in Australian army recruits, 1996. *Community Dentistry and Oral Epidemiology* 31(1):68–74.
- Hopcraft M & Morgan MV 2005. Dental caries experience in Australian Army recruits 2002–2003. *Australian Dental Journal* 50(1):16–20.
- Hopcraft MS & Morgan MV 2006. Pattern of dental caries experience on tooth surfaces in an adult population. *Community Dentistry and Oral Epidemiology* 34(3):174–83.
- Kingsford Smith D & Szuster F 2000. Aspects of tooth decay in recently arrived refugees. *Australian and New Zealand Journal of Public Health* 24(6):623–6.
- Loe H & Silness J 1963. Periodontal disease in pregnancy: 1. Prevalence and severity. *Acta Odontologica Scandinavica* 21:533–51.
- Marino R, Calache H, Wright C, Morgan M, Schofield SM & Minichiello V 2007. Profile of the oral health among ambulant older Greek and Italian migrants living in Melbourne. *Australian Dental Journal* 52(3):198–204.
- Marino R, Wright FA & Minas IH 2001. Oral health among Vietnamese using a community health centre in Richmond, Victoria. *Australian Dental Journal* 46(3):208–15.
- McGrath C & Bedi R 2002. Population based norming of the UK oral health related quality of life measure (OHQoL-UK). *British Dental Journal* 193:521–4.
- Morgan MV, Stonnill A & Laslett AM 1992. Dental caries amongst Royal Australian Navy recruits, 1988. *Australian Dental Journal* 37(3):201–4.
- NHANES (National Health and Nutrition Examination Survey). Dental examiners' procedures manual. Hyattsville, MD: US Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Health Statistics. Viewed 14 February 2007
<http://www.cdc.gov/nchs/data/nhanes/nhanes_03_04/DentalExaminers-2004.pdf>.
- Osborn M, Butler T & Barnard PD 2003. Oral health status of prison inmates – New South Wales, Australia. *Australian Dental Journal* 48(1):34–8.
- Saub R & Evans RW 2001. Dental needs of elderly hostel residents in inner Melbourne. *Australian Dental Journal* 46(3):198–202.
- Sheiham A, Steele JG, Marcenes W, Finch S & Walls AW 2002. The relationship between oral health status and body mass index among older people: a national survey of older people in Great Britain. *British Dental Journal* 192:703–6.
- Slade GD & Spencer AJ 1995. Periodontal attachment loss among adults aged 60+ in South Australia. *Community Dentistry and Oral Epidemiology* 23(4):237–42.

- Slade GD & Spencer AJ 1997. Distribution of coronal and root caries experience among persons aged 60+ in South Australia. *Australian Dental Journal* 42(3):178-84.
- Slade GD, Spencer AJ, Gorkic E & Andrews G 1993. Oral health status and treatment needs of non-institutionalized persons aged 60+ in Adelaide, South Australia. *Australian Dental Journal* 38(5):373-80.
- Slade GD, Spencer AJ & Roberts-Thomson KF (eds) 2007. *Australia's dental generations: The National Survey of Adult Oral Health 2004-06*. AIHW cat. no. DEN 165. Canberra: Australian Institute of Health and Welfare (Dental Statistics and Research Series No. 34).
- Smith K, Kruger E, Dyson K & Tennant M 2007. Oral health in rural and remote Western Australian indigenous communities: a two-year retrospective analysis of 999 people. *International Dental Journal* 57(2):93-9.
- Surgeon General 2000. *The health consequences of smoking: A report of the Surgeon General*. Atlanta, Georgia: U. S. Department of Health and Human Services, Centers for Disease Control and Prevention, National Center for Chronic Disease Prevention and Health Promotion, Office on Smoking and Health.
- Thomson WM, Slade GD & Spencer AJ 1995. Dental caries experience and use of prescription medications among people aged 60+ in South Australia. *Gerodontology* 12(12):104-10.
- Wright FA, Hammond RH & Lewis JM 1994. Changes in periodontal conditions of adults from Melbourne, Australia. *International Dental Journal* 44(3):207-14.

List of tables

Table 1: Distribution of examiners and examinations among states and territories	7
Table 2: Periods of data collection in states and territories	11
Table 3: Number and percentage of people sampled, interviewed and examined	12
Table 4: Percentage of people with selected sociodemographic and dental access characteristics in the NSW population and three age groups	15
Table 5: Percentage of adults with complete tooth loss	17
Table 6: Percentage of people with fewer than 21 teeth	19
Table 7: Percentage of dentate people who wear denture(s)	21
Table 8: Average number of teeth per person missing due to pathology	23
Table 9: Percentage of people with untreated coronal decay	25
Table 10: Percentage of people with untreated root decay	27
Table 11: Percentage of people with one or more filled teeth.....	29
Table 12: Average number of decayed, missing or filled teeth per person.....	31
Table 13: Percentage of people with moderate or severe periodontitis.....	33
Table 14: Percentage of people with 4+ mm periodontal pocket depth.....	35
Table 15: Percentage of people with 4+ mm clinical attachment loss.....	37
Table 16: Percentage of people with gingival inflammation.....	39
Table 17: Percentage of people visiting dentist within last 12 months.....	41
Table 18: Percentage of people who attended a private dental practice at last dental visit	43
Table 19: Percentage of people who paid for their last dental visit	45
Table 20: Percentage of people who received government-subsidised dental care in private sector	47
Table 21: Percentage of people who usually visit a dental professional at least once a year	49
Table 22: Percentage of people who have a dentist they usually attend.....	51
Table 23: Percentage of people who usually visit a dentist for a check-up.....	53
Table 24: Percentage of people who avoided or delayed dental care	55
Table 25: Percentage of people who reported that cost had prevented recommended dental treatment	57
Table 26: Percentage of people who would have a lot of difficulty paying a \$100 dental bill	59
Table 27: Percentage of people avoiding foods due to dental problems.....	61
Table 28: Percentage of people rating their oral health fair or poor	63
Table 29: Percentage of people experiencing toothache	65
Table 30: Percentage of people experiencing orofacial pain	67
Table 31: Percentage of people who need dentures	69
Table 32: Percentage of people who need an extraction or filling.....	71
Table 33: Percentage of people perceiving a need for check-up.....	73
Table 34: Percentage of people perceiving a need for treatment within 3 months	75
Table 35: Age-standardised comparison of health cardholders and non-health cardholders	77
Table 36: Age-standardised comparison of the dentally insured and the uninsured	79