

INCREASED *IN-VIVO* SENSITIVITY FOR TRANSGENE EXPRESSION IN MURINE NASAL AND LUNG AIRWAYS USING LUCIFERASE IMAGING

Patricia Cmielewski¹, Donald Anson², David Parsons^{1,3,4}

1. Respiratory and Sleep Medicine, Women's and Children's Hospital, Adelaide, South Australia

2. Gene Technology Unit, SA Pathology

3. Centre for Stem Cell Research, University of Adelaide

4. Women's and Children's Health Research Institute, Adelaide, South Australia

Introduction

Non-invasive bioluminescence imaging has allowed for rapid *in-vivo* quantification of long-lasting gene transfer in experimental animals. We are testing the longevity of a single nasal delivery of our lentiviral (LV) gene transfer system in mouse airways.

Methods

One nostril of C57Bl/6 mice was treated by a bolus instillation of a control (PBS) or the detergent lysophosphatidylcholine (LPC) one hour prior to delivery of a LV vector containing the reporter-gene luciferase (Luc) at 1.8×10^{10} tu/ml. Imaging to detect luminescence was via the IVIS system (Xenogen) 10-15 minutes after a 50 μ l intranasal bolus of the substrate D-luciferin (15mg/ml PBS stock), at 1 week (Fig. 1a) and 1, 3 and 6 months (Fig. 1b) post LV.

Results

LPC pre-treated LV gene transfer resulted in significantly greater nasal gene transfer compared to PBS pre-treatment at all time points (* $p < 0.05$, ANOVA). A statistically significant reduction in nasal luminescence was noted at 3 and 6 months compared to 1 week for LPC pretreated animals (# $p < 0.05$, RM ANOVA, Fig. 2). Luciferase activity was also detected in the lung in both groups of mice (Fig. 3). At the 6 month time point an increase in lung luminescence was observed in mice pre-treated with PBS prior to LV (# $p < 0.05$, RM ANOVA).

Results




Fig. 1a. 1 week LV-luciferase luminescence PBS (left) vs LPC (right)




Fig. 1b. 6 months LV-luciferase luminescence; same animals as above




Fig. 2. Nasal LV-luciferase luminescence




Fig. 3. Lung LV-luciferase luminescence

Conclusion

Nasal lentiviral luciferase gene expression persisted for at least 6 months after a single dose, and was enhanced with LPC pre-treatment. The luciferase gene expression could also be detected in the lung, with or without LPC pre-treatment. This is in contrast to LacZ reporter gene expression; after similar nasal administration LacZ was only detected in nasal airways. The luciferase reporter gene has greater sensitivity for airway gene transfer detection.

Acknowledgements

NH&MRC and philanthropic donors (www.Cure4CF.org).